

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/47, A61K 38/16, C12N 15/62, C07K 16/18		A1	(11) International Publication Number: WO 98/50552 (43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/US98/09584 (22) International Filing Date: 6 May 1998 (06.05.98) (30) Priority Data: 60/045,703 6 May 1997 (06.05.97) US (71) Applicant: ZYMOGENETICS, INC. [US/US]; 1201 Eastlake Avenue East, Seattle, WA 98102 (US). (72) Inventors: SHEPPARD, Paul, O.; 20717 N.E. 2nd Street, Redmond, WA 98053 (US). GROSSMANN, Angelika; 2809 N.E. 60th, Seattle, WA 98115 (US). (74) Agent: SAWISLAK, Deborah, A.; ZymoGenetics, Inc., 1201 Eastlake Avenue East, Seattle, WA 98102 (US).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NOVEL TUMOR ANTIGENS			
(57) Abstract The present invention relates to polynucleotide and polypeptide molecules for a secreted polypeptide designated zsig15. The polypeptides, and polynucleotides encoding them, may be used for mapping chromosome 19 and markers for tumor growth. The present invention also includes antibodies to the zsig15 polypeptides.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Description

5

NOVEL TUMOR ANTIGENS

BACKGROUND OF THE INVENTION

Prostate, colon and breast tissues are all
10 associated with a high prevalence of cancer, and are in
fact with lung cancer, the most deadly forms of cancer in
the United States today.

Prostate cancer is the most prevalent form of
cancer in men in the United States. It is anticipated
15 that an estimated 334,500 new cases of prostate cancer
will be diagnosed in 1997 (in Cancer Facts and Figures:
1997, American Cancer Society, Atlanta, 1997). The
incidence rate for prostate cancer has increased 50%
between 1989 and 1993

20 Early diagnosis is critical for survival, for
example, if a tumor is discovered while it is still
localized, the 5 year survival rate is 99%. However,
early diagnosis at the present time is limited to
relatively unreliable methods. These methods include a
25 digital rectal exam and prostate specific antigen (PSA)
test. However, digital exam may require that the tumor
has become quite large and the cancer may be at a more
advanced stage. Ultrasonography has demonstrated that 20%
to 30% of cancers detected using this method are not
30 detectable using PSA tests. In addition, false positives
are a significant problem as well (in Cancer: Principles
and Practice of Oncology, DeVita, Hellman and Rosenberg
(eds), J.B. Lippincott Company, Philadelphia, pp:538-589,
1993).

35 Breast cancer is the second major cause of
cancer death in women. Approximately 110/100,000 women
will be diagnosed with breast cancer each year, and it is

expected that in 1997 there will be 43,900 deaths related to breast cancer (in Cancer Facts and Figures: 1997, American Cancer Society, Atlanta, 1997). Early detection is the key to survival and mammograms are generally
5 accepted as the most valuable tool that is available for early detection. Once a tumor is recognizable by physical symptoms the tumor has generally progressed.

An estimated 94,100 cases of colon cancer will be identified in 1997, and 10% of all cancer deaths will
10 be related to colorectal cancer (in Cancer Facts and Figures: 1997, American Cancer Society, Atlanta, 1997). While diagnosis and treatment have been steadily improving, early detection is important for survival. In addition, the cells in these tissues have a high rate
15 of turnover and metabolic activity, and therefore, generally produce a large spectrum of proteins. Included are proteins with secretory functions, and this is particularly common in tissue involved in transporting substances to the outside of the body. One type of
20 substance produced by secretory tissues is anti-microbial agents, in particular antimicrobial proteins. These proteins can act via several mechanisms that include: enzymatic mechanisms, such as, breaking down microbial protective substances or the microbial cell wall/membrane
25 or the proteins can be very sticky and trap microorganisms. The antimicrobial proteins produced by secretory tissue play an important role in maintaining the body's ability to prevent infection in a more non-specific way than is generally associated with components of the
30 immune system.

Thus, the present invention provides proteins that will be valuable as markers for changes in metabolism highly susceptible to cancerous growth, as well as other uses that should be apparent to those skilled in the art
35 from the teachings herein.

SUMMARY OF THE INVENTION

The present invention provides an isolated polynucleotide molecule selected from the group consisting of: a) polynucleotide molecules comprising a nucleotide
5 sequence as shown in SEQ ID NO: 1 from nucleotide 97 to nucleotide 1215; b) orthologs of (a); (c) allelic variants of (a) or (b); d) polynucleotide molecules that encode a polypeptide that is at least 80% identical to the amino acid sequence of SEQ ID NO: 2 from amino acid residue 22
10 (Leu) to amino acid residue 394 (Glu); and e) degenerate nucleotide sequences of (a), (b), (c) or (d).

In other embodiments, the polynucleotide molecule is selected from nucleotide sequences as shown in SEQ ID NO: 1 from nucleotides 97-1344, 79-1344, 34-1344,
15 97-582, 65-1215, or 65-1344; and the corresponding polynucleotide molecules that encode a polypeptide that is at least 80% identical to the amino acid sequence of SEQ ID NO: 2 from residues 22-437, 16-437, or 1-347, respectively.

20 In aspects, the present invention provides isolated polynucleotide molecules comprising a sequence of nucleotides from nucleotide 97-582, 655-1215, or 655-1344 of SEQ ID NO: 1.

In another aspect, the present invention
25 provides a isolated polynucleotide molecule encoding a fusion protein comprising a first polypeptide and a second polypeptide, said first polypeptide encoded by a sequence of nucleotides as shown in SEQ ID NO: 1 from nucleotide 97 to nucleotide 582, and a second polypeptide encoded by a
30 sequence of nucleotides as shown in SEQ ID NO: 1 from nucleotide 655 to nucleotide 1215.

In another aspect, the present invention provides an expression vector comprising the following operably linked elements: a transcription promoter; a DNA
35 segment selected from the group consisting of: (a) polynucleotide molecules comprising a nucleotide sequence

as shown in SEQ ID NO: 1 from nucleotide 97 to nucleotide 1215; (b) orthologs of (a); (c) allelic variants of (a) or (b); (d) polynucleotide molecules that encode a polypeptide that is at least 80% identical to the amino acid sequence of SEQ ID NO: 2 from amino acid residue 22 (Leu) to amino acid residue 394 (Glu); and (e) degenerate nucleotide sequences of (a), (b), (c) or (d); and a transcription terminator.

In another aspect, the present invention provides a cultured cell into which has been introduced an expression vector according as described above, wherein said cell expresses the polypeptide encoded by the DNA segment.

In another aspect, the present invention provides an isolated polypeptide selected from the group consisting of: (a) polypeptide molecules comprising an amino acid sequence as shown in SEQ ID NO: 2 from amino acid residue 22 (Leu) to amino acid residue 394 (Glu); (b) orthologs of (a); (c) allelic variants of (a) or (b); and (d) polypeptide molecules that are at least 80% identical to the amino acids of SEQ ID NO: 2 from amino acid residue 22 (Leu) to amino acid residue 394 (Glu).

In other embodiments, the present invention provides isolated polypeptides wherein said polypeptide molecules comprise an amino acid sequence as shown in SEQ ID NO: 2 from residues 22-437, 16-437 and 1-437.

In another aspect, the present invention provides isolated polypeptides comprising a sequence of amino acids as shown in SEQ ID NO: 2 from residue 22-183, 208-394 and 208-437.

In another aspect, the present invention provides an isolated fusion protein comprising a first polypeptide and a second polypeptide, said first polypeptide comprising a sequence of amino acid residues as shown in SEQ ID NO: 2 from residues 22-183, and a

second polypeptide comprising a sequence amino acid residues as shown in SEQ ID NO: 2 from residues 208-394.

In another embodiment, the present invention provides an isolated polypeptide that is additionally
5 covalently linked at the N-terminus or C-terminus to a moiety selected from the group consisting of affinity tags, toxins, radionucleotides, enzymes or fluorophores.

In another aspect, the present invention provides antibodies that specifically bind to an epitope
10 of a polypeptide comprising a sequence of amino acids as shown in SEQ ID NO: 2 from residue 1-437.

In another aspect, the present invention provides a method of producing a polypeptide comprising:
culturing a cell into which has been introduced an
15 expression vector as described above, whereby said cell expresses a polypeptide encoded by the DNA segment; and recovering the polypeptide.

In another aspect, the present invention provides an oligonucleotide probe or primer comprising at
20 least 14 contiguous nucleotides of polynucleotide of SEQ ID NO: 1 from nucleotide 34 to nucleotide 1344.

DETAILED DESCRIPTION OF THE INVENTION

25 Prior to describing the present invention in detail, it may be helpful to define certain terms used herein:

The term "affinity tag" is used herein to denote a polypeptide segment that can be attached to a second
30 polypeptide to provide for purification or detection of the second polypeptide or provide sites for attachment of the second polypeptide to a substrate. In principal, any peptide or protein for which an antibody or other specific binding agent is available can be used as an affinity tag.
35 Affinity tags include a poly-histidine tract, protein A (Nilsson et al., EMBO J. 4:1075, 1985; Nilsson et al.,

Methods Enzymol. 198:3, 1991), glutathione S transferase (Smith and Johnson, Gene 67:31, 1988), Glu-Glu affinity tag (Grussenmeyer et al., Proc. Natl. Acad. Sci. USA 82:7952-4, 1985), substance P, FlagTM peptide (Hopp et al., 5 Biotechnology 6:1204-10, 1988), streptavidin binding peptide, or other antigenic epitope or binding domain. See, in general, Ford et al., Protein Expression and Purification 2: 95-107, 1991. DNAs encoding affinity tags are available from commercial suppliers (e.g., Pharmacia 10 Biotech, Piscataway, NJ).

The term "allelic variant" is used herein to denote any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in 15 phenotypic polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequence. The term allelic variant is also used herein to denote a protein encoded by an allelic 20 variant of a gene.

The terms "amino-terminal" and "carboxyl-terminal" are used herein to denote positions within polypeptides. Where the context allows, these terms are used with reference to a particular sequence or portion of 25 a polypeptide to denote proximity or relative position. For example, a certain sequence positioned carboxyl-terminal to a reference sequence within a polypeptide is located proximal to the carboxyl terminus of the reference sequence, but is not necessarily at the carboxyl terminus 30 of the complete polypeptide.

The term "complement/anti-complement pair" denotes non-identical moieties that form a non-covalently associated, stable pair under appropriate conditions. For instance, biotin and avidin (or streptavidin) are 35 prototypical members of a complement/anti-complement pair. Other exemplary complement/anti-complement pairs include

receptor/ligand pairs, antibody/antigen (or hapten or epitope) pairs, sense/antisense polynucleotide pairs, and the like. Where subsequent dissociation of the complement/anti-complement pair is desirable, the
5 complement/anti-complement pair preferably has a binding affinity of $<10^9 \text{ M}^{-1}$.

The term "complements of a polynucleotide molecule" is a polynucleotide molecule having a complementary base sequence and reverse orientation as
10 compared to a reference sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

The term "contig" denotes a polynucleotide that has a contiguous stretch of identical or complementary
15 sequence to another polynucleotide. Contiguous sequences are said to "overlap" a given stretch of polynucleotide sequence either in their entirety or along a partial stretch of the polynucleotide. For example, representative contigs to the polynucleotide sequence 5'-
20 ATGGCTTAGCTT-3' are 5'-TAGCTTgagtct-3' and 3'-gtcgactACCGA-5'.

The term "degenerate nucleotide sequence" denotes a sequence of nucleotides that includes one or more degenerate codons (as compared to a reference
25 polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "expression vector" is used to denote a
30 DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments include promoter and terminator sequences, and may also include one or more origins of
35 replication, one or more selectable markers, an enhancer, a polyadenylation signal, etc. Expression vectors are

generally derived from plasmid or viral DNA, or may contain elements of both.

The term "isolated", when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, Nature 316:774-78, 1985).

An "isolated" polypeptide or protein is a polypeptide or protein that is found in a condition other than its native environment, such as apart from blood and animal tissue. In a preferred form, the isolated polypeptide is substantially free of other polypeptides, particularly other polypeptides of animal origin. It is preferred to provide the polypeptides in a highly purified form, i.e., greater than 95% pure, more preferably greater than 99% pure. When used in this context, the term "isolated" does not exclude the presence of the same polypeptide in alternative physical forms, such as dimers or alternatively glycosylated or derivatized forms.

The term "operably linked", when referring to DNA segments, indicates that the segments are arranged so that they function in concert for their intended purposes, e.g., transcription initiates in the promoter and proceeds through the coding segment to the terminator.

The term "ortholog" denotes a polypeptide or protein obtained from one species that is the functional

counterpart of a polypeptide or protein from a different species. Sequence differences among orthologs are the result of speciation.

"Paralogs" are distinct but structurally related proteins made by an organism. Paralogs are believed to arise through gene duplication. For example, α -globin, β -globin, and myoglobin are paralogs of each other.

A "polynucleotide" is a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic molecules. Sizes of polynucleotides are expressed as base pairs (abbreviated "bp"), nucleotides ("nt"), or kilobases ("kb"). Where the context allows, the latter two terms may describe polynucleotides that are single-stranded or double-stranded. When the term is applied to double-stranded molecules it is used to denote overall length and will be understood to be equivalent to the term "base pairs". It will be recognized by those skilled in the art that the two strands of a double-stranded polynucleotide may differ slightly in length and that the ends thereof may be staggered as a result of enzymatic cleavage; thus all nucleotides within a double-stranded polynucleotide molecule may not be paired. Such unpaired ends will in general not exceed 20 nt in length.

A "polypeptide" is a polymer of amino acid residues joined by peptide bonds, whether produced naturally or synthetically. Polypeptides of less than about 10 amino acid residues are commonly referred to as "peptides".

The term "promoter" is used herein for its art-recognized meaning to denote a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter

sequences are commonly, but not always, found in the 5' non-coding regions of genes.

The term "receptor" denotes a cell-associated protein that binds to a bioactive molecule (i.e., a ligand) and mediates the effect of the ligand on the cell. Membrane-bound receptors are characterized by a multi-domain structure comprising an extracellular ligand-binding domain and an intracellular effector domain that is typically involved in signal transduction. Binding of ligand to receptor results in a conformational change in the receptor that causes an interaction between the effector domain and other molecule(s) in the cell. This interaction in turn leads to an alteration in the metabolism of the cell. Metabolic events that are linked to receptor-ligand interactions include gene transcription, phosphorylation, dephosphorylation, increases in cyclic AMP production, mobilization of cellular calcium, mobilization of membrane lipids, cell adhesion, hydrolysis of inositol lipids and hydrolysis of phospholipids. Most nuclear receptors also exhibit a multi-domain structure, including an amino-terminal, transactivating domain, a DNA binding domain and a ligand binding domain. In general, receptors can be membrane bound, cytosolic or nuclear; monomeric (e.g., thyroid stimulating hormone receptor, beta-adrenergic receptor) or multimeric (e.g., PDGF receptor, growth hormone receptor, IL-3 receptor, GM-CSF receptor, G-CSF receptor, erythropoietin receptor and IL-6 receptor).

The term "secretory signal sequence" denotes a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

The term "splice variant" is used herein to denote alternative forms of RNA transcribed from a gene. Splice variation arises naturally through use of alternative splicing sites within a transcribed RNA molecule, or less commonly between separately transcribed RNA molecules, and may result in several mRNAs transcribed from the same gene. Splice variants may encode polypeptides having latered amino acid sequence. The term splice variant is also used herein to denote a protein encoded by a splice variant of an mRNA transcribed from a gene.

Molecular weights and lengths of polymers determined by imprecise analytical methods (e.g., gel electrophoresis) will be understood to be approximate values. When such a value is expressed as "about" X or "approximately" X, the stated value of X will be understood to be accurate to $\pm 10\%$.

All references cited herein are incorporated by reference in their entirety.

20

The present invention is based in part upon the discovery of a novel DNA sequence that encodes a secreted polypeptide. Analysis of the tissue distribution of the mRNA corresponding to this novel DNA revealed that expression was highest in prostate and bone marrow, followed by apparent but decreased expression levels in spinal cord, colon and pancreas. Transcripts of 1.7 and 2.5 kb in size were identified by northern analysis. The polypeptide has been designated zsig15. Northern blots of tumor and normal tissue further revealed mRNA expression in colon and breast at various stages of tumor development and/or normal tissue.

The novel zsig15 polypeptides of the present invention were initially identified by querying an EST database for sequences possessing a putative secretion signal. An EST sequence was discovered and was

subsequently mapped to chromosome 19;q13.1-913.2. The EST sequence was originally derived from a colon tumor library.

The nucleotide sequence of zsig15 is described in SEQ ID NO. 1, and its deduced amino acid sequence is described in SEQ ID NO. 2. Analysis of the DNA encoding a zsig15 polypeptide (SEQ ID NO: 1) revealed an open reading frame encoding 437 amino acids (SEQ ID NO: 2) comprising a signal peptide of from 15 to 21 amino acid residues (residue 1 to residues 15 to 21 of SEQ ID NO: 2) and comprising a mature polypeptide of from 416 to 422 amino acids (residues 15-21 to residue 437 of SEQ ID NO: 2).

The polypeptide of the present invention contains at least 4 distinct regions. Two of these regions are direct repeats of each other. Starting from the mature peptide amino terminus (residue 16 or 22) region 1 is the first repeated domain as shown in SEQ ID NO: 2 from amino acid residue 22 (Leu) to residue 183 (Pro) and contains 12 Cysteine residues. Region 2 (residues 184 (Gly) to residue 207 (Asp) of SEQ ID NO: 2) appears to be a linker region and contains a pair of Cysteines. Region 3 (amino acid residue 208 (Phe) to residue 376 (Pro) of SEQ ID NO: 2) is the second repeated domain, and contains 12 Cysteines. Both repeated domains are predicted to be all beta-strand structure, which suggests a tertiary structure similarity with IG domains and cytokine receptor domains. The fourth domain is the C-terminus of the polypeptide (residues 377-437 of SEQ ID NO: 2) and contains a hydrophobic stretch which is a possible anchor for dimerization, membrane association, or hydrophobic matrix binding for the polypeptide, and contains a pair of Cysteines. The Arg-Lys, as shown in SEQ ID NO: 2 at residues 205-206 is a common processing site. Another putative processing site is found at residues 395-396 (Lys-Arg) of SEQ ID NO: 2. This cleavage site may serve to release the polypeptide as a free

secretory component from the membrane-bound form of the protein and make it soluble. Generation of a free secretory component from a membrane-bound protein has been suggested for TNF- α (Rosendahl et al., J. Biol. Chem. 5 272(39):24588-24593, 1997).

The present invention provides polynucleotide molecules, including DNA and RNA molecules, that encode the zsig15 polypeptides disclosed herein. Those skilled in the art will readily recognize that, in view of the
10 degeneracy of the genetic code, considerable sequence variation is possible among these polynucleotide molecules. SEQ ID NO:4 is a degenerate DNA sequence that encompasses all DNAs that encode the zsig15 polypeptide of SEQ ID NO:2. Those skilled in the art will recognize that
15 the degenerate sequence of SEQ ID NO:4 also provides all RNA sequences encoding SEQ ID NO:2 by substituting U for T. Thus, zsig15 polypeptide-encoding polynucleotides comprising nucleotide 1 to nucleotide 1311 of SEQ ID NO:4 and their RNA equivalents are contemplated by the present
20 invention. Table 1 sets forth the one-letter codes used within SEQ ID NO:4 to denote degenerate nucleotide positions. "Resolutions" are the nucleotides denoted by a code letter. "Complement" indicates the code for the complementary nucleotide(s). For example, the code Y
25 denotes either C or T, and its complement R denotes A or G, A being complementary to T, and G being complementary to C.

TABLE 1

Nucleotide	Resolution	Nucleotide	Complement
A	A	T	T
C	C	G	G
G	G	C	C
T	T	A	A
R	A G	Y	C T
Y	C T	R	A G
M	A C	K	G T
K	G T	M	A C
S	C G	S	C G
W	A T	W	A T
H	A C T	D	A G T
B	C G T	V	A C G
V	A C G	B	C G T
D	A G T	H	A C T
N	A C G T	N	A C G T

The degenerate codons used in SEQ ID NO:4,
5 encompassing all possible codons for a given amino acid,
are set forth in Table 2.

TABLE 2

Amino Acid	One Letter Code	Codons	Degenerate Codon
Cys	C	TGC TGT	TGY
Ser	S	AGC AGT TCA TCC TCG TCT	WSN
Thr	T	ACA ACC ACG ACT	ACN
Pro	P	CCA CCC CCG CCT	CCN
Ala	A	GCA GCC GCG GCT	GCN
Gly	G	GGA GGC GGG GGT	GGN
Asn	N	AAC AAT	AAY
Asp	D	GAC GAT	GAY
Glu	E	GAA GAG	GAR
Gln	Q	CAA CAG	CAR
His	H	CAC CAT	CAY
Arg	R	AGA AGG CGA CGC CGG CGT	MGN
Lys	K	AAA AAG	AAR
Met	M	ATG	ATG
Ile	I	ATA ATC ATT	ATH
Leu	L	CTA CTC CTG CTT TTA TTG	YTN
Val	V	GTA GTC GTG GTT	GTN
Phe	F	TTC TTT	TTY
Tyr	Y	TAC TAT	TAY
Trp	W	TGG	TGG
Ter	.	TAA TAG TGA	TRR
Asn Asp	B		RAY
Glu Gln	Z		SAR
Any	X		NNN

One of ordinary skill in the art will appreciate that some ambiguity is introduced in determining a degenerate codon, representative of all possible codons encoding each amino acid. For example, the degenerate
5 codon for serine (WSN) can, in some circumstances, encode arginine (AGR), and the degenerate codon for arginine (MGN) can, in some circumstances, encode serine (AGY). A similar relationship exists between codons encoding phenylalanine and leucine. Thus, some polynucleotides
10 encompassed by the degenerate sequence may encode variant amino acid sequences, but one of ordinary skill in the art can easily identify such variant sequences by reference to the amino acid sequence of SEQ ID NO:2. Variant sequences can be readily tested for functionality as described
15 herein.

One of ordinary skill in the art will also appreciate that different species can exhibit "preferential codon usage." In general, see, Grantham, et al., Nuc. Acids Res. 8:1893-912, 1980; Haas, et al. Curr.
20 Biol. 6:315-24, 1996; Wain-Hobson, et al., Gene 13:355-64, 1981; Grosjean and Fiers, Gene 18:199-209, 1982; Holm, Nuc. Acids Res. 14:3075-87, 1986; Ikemura, J. Mol. Biol. 158:573-97, 1982. As used herein, the term "preferential codon usage" or "preferential codons" is a term of art
25 referring to protein translation codons that are most frequently used in cells of a certain species, thus favoring one or a few representatives of the possible codons encoding each amino acid (See Table 2). For example, the amino acid Threonine (Thr) may be encoded by
30 ACA, ACC, ACG, or ACT, but in mammalian cells ACC is the most commonly used codon; in other species, for example, insect cells, yeast, viruses or bacteria, different Thr codons may be preferential. Preferential codons for a particular species can be introduced into the
35 polynucleotides of the present invention by a variety of methods known in the art. Introduction of preferential codon sequences into recombinant DNA can, for example,

enhance production of the protein by making protein translation more efficient within a particular cell type or species. Therefore, the degenerate codon sequence disclosed in SEQ ID NO:4 serves as a template for
5 optimizing expression of polynucleotides in various cell types and species commonly used in the art and disclosed herein. Sequences containing preferential codons can be tested and optimized for expression in various species, and tested for functionality as disclosed herein.

10 Within preferred embodiments of the invention the isolated polynucleotides will hybridize to similar sized regions of SEQ ID NO:1, or a sequence complementary thereto, under stringent conditions. In general, stringent conditions are selected to be about 5°C lower
15 than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typical stringent conditions are those in
20 which the salt concentration is up to about 0.03 M at pH 7 and the temperature is at least about 60°C.

As previously noted, the isolated polynucleotides of the present invention include DNA and RNA. Methods for preparing DNA and RNA are well known in
25 the art. In general, RNA is isolated from a tissue or cell that produces large amounts of zsig15 RNA. Such tissues and cells are identified by Northern blotting (Thomas, Proc. Natl. Acad. Sci. USA 77:5201, 1980), and include prostate, bone marrow, colon, breast and spinal
30 cord. Total RNA can be prepared using guanidine HCl extraction followed by isolation by centrifugation in a CsCl gradient (Chirgwin et al., Biochemistry 18:52-94, 1979). Poly (A)⁺ RNA is prepared from total RNA using the method of Aviv and Leder (Proc. Natl. Acad. Sci. USA
35 69:1408-12, 1972). Complementary DNA (cDNA) is prepared

from poly(A)⁺ RNA using known methods. In the alternative, genomic DNA can be isolated. Polynucleotides encoding zsig15 polypeptides are then identified and isolated by, for example, hybridization or PCR.

5 A full-length clone encoding zsig15 can be obtained by conventional cloning procedures. Complementary DNA (cDNA) clones are preferred, although for some applications (e.g., expression in transgenic animals) it may be preferable to use a genomic clone, or
10 to modify a cDNA clone to include at least one genomic intron. Methods for preparing cDNA and genomic clones are well known and within the level of ordinary skill in the art, and include the use of the sequence disclosed herein, or parts thereof, for probing or priming a library.
15 Expression libraries can be probed with antibodies to zsig15, receptor fragments, or other specific binding partners.

 The present invention further provides counterpart polypeptides and polynucleotides from other
20 species (orthologs). These species include, but are not limited to mammalian, avian, amphibian, reptile, fish, insect and other vertebrate and invertebrate species. Of particular interest are zsig15 polypeptides from other mammalian species, including murine, porcine, ovine,
25 bovine, canine, feline, equine, and other primate polypeptides. Orthologs of human zsig15 can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a cDNA can be cloned using mRNA
30 obtained from a tissue or cell type that expresses zsig15 as disclosed herein. Suitable sources of mRNA can be identified by probing Northern blots with probes designed from the sequences disclosed herein. A library is then prepared from mRNA of a positive tissue or cell line. A
35 zsig15-encoding cDNA can then be isolated by a variety of methods, such as by probing with a complete or partial

human cDNA or with one or more sets of degenerate probes based on the disclosed sequences. A cDNA can also be cloned using the polymerase chain reaction, or PCR (Mullis, U.S. Patent No. 4,683,202), using primers
5 designed from the representative human zsig15 sequences disclosed herein. Within an additional method, the cDNA library can be used to transform or transfect host cells, and expression of the cDNA of interest can be detected with an antibody to zsig15 polypeptide. Similar techniques
10 can also be applied to the isolation of genomic clones.

Those skilled in the art will recognize that the sequence disclosed in SEQ ID NO:1 represents a single allele of human zsig15 and that allelic variation and alternative splicing are expected to occur. Allelic
15 variants can be cloned by probing cDNA or genomic libraries from different individuals according to standard procedures. Allelic variants of the DNA sequence shown in SEQ ID NO: 1, including those containing silent mutations and those in which mutations result in amino
20 acid sequence changes, are within the scope of the present invention, as are proteins which are allelic variants of SEQ ID NO:2. cDNAs generated from alternatively spliced mRNAs, which retain the properties of the zsig15 polypeptide are included within the scope of the present
25 invention, as are polypeptides encoded by such cDNAs and mRNAs. Allelic variants and splice variants of these sequences can be cloned by probing cDNA or genomic libraries from different individuals or tissues according to standard procedures known in the art.

30 The highly conserved amino acids in zsig15 can be used as a tool to identify new family members. For instance, reverse transcription-polymerase chain reaction (RT-PCR) can be used to amplify sequences encoding the conserved regions from RNA obtained from a variety of
35 tissue sources or cell lines. In particular, highly

degenerate primers designed from the zsig15 sequences are useful for this purpose.

The present invention also provides isolated zsig15 polypeptides that are substantially homologous to the polypeptides of SEQ ID NO:2 and their orthologs. The term "substantially homologous" is used herein to denote polypeptides having 60%, preferably 80%, more preferably at least 90%, and most preferably at least 95%, sequence identity to the sequences shown in SEQ ID NO:2 or their orthologs. Percent sequence identity is determined by conventional methods. See, for example, Altschul et al., Bull. Math. Bio. 48: 603-616, 1986 and Henikoff and Henikoff, Proc. Natl. Acad. Sci. USA 89:10915-10919, 1992. Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "blosum 62" scoring matrix of Henikoff and Henikoff (ibid.) as shown in Table 3 (amino acids are indicated by the standard one-letter codes).

21

Table 3

	A	R	N	D	C	Q	E	G	H	I	L	K	M	F	P	S	T	W	Y	V	
	A	4																			
5	R	-1	5																		
	N	-2	0	6																	
	D	-2	-2	1	6																
	C	0	-3	-3	-3	9															
	Q	-1	1	0	0	-3	5														
10	E	-1	0	0	2	-4	2	5													
	G	0	-2	0	-1	-3	-2	-2	6												
	H	-2	0	1	-1	-3	0	0	-2	8											
	I	-1	-3	-3	-3	-1	-3	-3	-4	-3	4										
	L	-1	-2	-3	-4	-1	-2	-3	-4	-3	2	4									
15	K	-1	2	0	-1	-3	1	1	-2	-1	-3	-2	5								
	M	-1	-1	-2	-3	-1	0	-2	-3	-2	1	2	-1	5							
	F	-2	-3	-3	-3	-2	-3	-3	-3	-1	0	0	-3	0	6						
	P	-1	-2	-2	-1	-3	-1	-1	-2	-2	-3	-3	-1	-2	-4	7					
	S	1	-1	1	0	-1	0	0	0	-1	-2	-2	0	-1	-2	-1	4				
20	T	0	-1	0	-1	-1	-1	-1	-2	-2	-1	-1	-1	-1	-2	-1	1	5			
	W	-3	-3	-4	-4	-2	-2	-3	-2	-2	-3	-2	-3	-1	1	-4	-3	-2	11		
	Y	-2	-2	-2	-3	-2	-1	-2	-3	2	-1	-1	-2	-1	3	-3	-2	-2	2	7	
	V	0	-3	-3	-3	-1	-2	-2	-3	-3	3	1	-2	1	-1	-2	-2	0	-3	-1	4

The percent identity is then calculated as:
Total number of identical matches

5
$$\frac{\text{[length of the longer sequence plus the number of gaps introduced into the longer sequence in order to align the two sequences]}}{\text{[length of the longer sequence plus the number of gaps introduced into the longer sequence in order to align the two sequences]}} \times 100$$

Sequence identity of polynucleotide molecules is determined by similar methods using a ratio as disclosed above.

10 Variant zsig15 polypeptides or substantially homologous zsig15 polypeptides are characterized as having one or more amino acid substitutions, deletions or additions. These changes are preferably of a minor nature, that is conservative amino acid substitutions
15 (see Table 4) and other substitutions that do not significantly affect the folding or activity of the polypeptide; small deletions, typically of one to about 30 amino acids; and small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue,
20 a small linker peptide of up to about 20-25 residues, or an affinity tag.

As described previously, the polypeptides of the present invention have at least 4 distinct regions. These regions (residues 22-183, residues 184-207,
25 residues 208-376 and residues 377-437 of SEQ ID NO: 2) can be used independently or as fusion proteins in combination with other regions from polypeptides of the present invention or elsewhere. The present invention thus includes polypeptides of from 61 to 437 amino acid
30 residues that comprise a sequence that is at least 80%, preferably at least 90%, and more preferably 95% or more identical to the corresponding region of SEQ ID NO:2. When combining polypeptides (or the polynucleotides that encode them), in one aspect of the present invention, the
35 fusion protein comprises a first polypeptide and a second polypeptide, said first polypeptide comprises amino acid

residues 22 (Leu) to 183 (Pro), or the polynucleotides encoding said polypeptide, and said second polypeptide comprises amino acid residues 208 (Phe) to 394 (Glu) or the polynucleotides encoding said polypeptide.

5 Polypeptides comprising covalently linked moieties, such as, affinity tags can further comprise a proteolytic cleavage site between the zsig15 polypeptide and the affinity tag. Preferred such sites include thrombin cleavage sites and factor Xa cleavage sites.

10 Table 4

Conservative amino acid substitutions

15	Basic:	arginine
		lysine
		histidine
	Acidic:	glutamic acid
		aspartic acid
20	Polar:	glutamine
		asparagine

Table 4 cont.

25	Hydrophobic:	leucine
		isoleucine
		valine
	Aromatic:	phenylalanine
		tryptophan
		tyrosine
30	Small:	glycine
		alanine
		serine
		threonine
		methionine

35 The present invention further provides a variety of other polypeptide fusions and related

multimeric proteins comprising one or more polypeptide fusions. For example, a zsig15 polypeptide can be prepared as a fusion to a dimerizing protein as disclosed in U.S. Patents Nos. 5,155,027 and 5,567,584. Preferred
5 dimerizing proteins in this regard include immunoglobulin constant region domains. Immunoglobulin-zsig15 polypeptide fusions can be expressed in genetically engineered cells to produce a variety of multimeric zsig15 analogs. Auxiliary domains can be fused to zsig15
10 polypeptides to target them to specific cells, tissues, or macromolecules (e.g., collagen). For example, a zsig15 polypeptide or protein could be targeted to a predetermined cell type by fusing a zsig15 polypeptide to a ligand that specifically binds to a receptor on the
15 surface of the target cell. In this way, polypeptides and proteins can be targeted for therapeutic or diagnostic purposes. A zsig15 polypeptide can be fused to two or more moieties, such as an affinity tag for purification and a targeting domain. Polypeptide fusions
20 can also comprise one or more cleavage sites, particularly between domains. Fusion with moieties may be either at the N-terminus or C-terminus of the zsig15 polypeptide. See, Tuan et al., Connective Tissue Research 34:1-9, 1996.

25 The proteins of the present invention can also comprise non-naturally occurring amino acid residues. Non-naturally occurring amino acids include, without limitation, trans-3-methylproline, 2,4-methanoproline, cis-4-hydroxyproline, trans-4-hydroxyproline, N-
30 methylglycine, allo-threonine, methylthreonine, hydroxyethylcysteine, hydroxyethylhomocysteine, nitroglutamine, homoglutamine, pipecolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, 3,3-dimethylproline, tert-leucine,
35 norvaline, 2-azaphenylalanine, 3-azaphenylalanine, 4-

azaphenylalanine, and 4-fluorophenylalanine. Several methods are known in the art for incorporating non-naturally occurring amino acid residues into proteins. For example, an *in vitro* system can be employed wherein

5 nonsense mutations are suppressed using chemically aminoacylated suppressor tRNAs. Methods for synthesizing amino acids and aminoacylating tRNA are known in the art. Transcription and translation of plasmids containing nonsense mutations is carried out in a cell-free system

10 comprising an *E. coli* S30 extract and commercially available enzymes and other reagents. Proteins are purified by chromatography. See, for example, Robertson et al., J. Am. Chem. Soc. 113:2722, 1991; Ellman et al., Methods Enzymol. 202:301, 1991; Chung et al., Science

15 259:806-809, 1993; and Chung et al., Proc. Natl. Acad. Sci. USA 90:10145-10149, 1993). In a second method, translation is carried out in *Xenopus* oocytes by microinjection of mutated mRNA and chemically aminoacylated suppressor tRNAs (Turcatti et al., J. Biol.

20 Chem. 271:19991-19998, 1996). Within a third method, *E. coli* cells are cultured in the absence of a natural amino acid that is to be replaced (e.g., phenylalanine) and in the presence of the desired non-naturally occurring amino acid(s) (e.g., 2-azaphenylalanine, 3-azaphenylalanine, 4-

25 azaphenylalanine, or 4-fluorophenylalanine). The non-naturally occurring amino acid is incorporated into the protein in place of its natural counterpart. See, Koide et al., Biochem. 33:7470-7476, 1994. Naturally occurring amino acid residues can be converted to non-naturally

30 occurring species by *in vitro* chemical modification. Chemical modification can be combined with site-directed mutagenesis to further expand the range of substitutions (Wynn and Richards, Protein Sci. 2:395-403, 1993).

A limited number of non-conservative amino

35 acids, amino acids that are not encoded by the genetic

code, non-naturally occurring amino acids, and unnatural amino acids may be substituted for zsig15 amino acid residues.

Essential amino acids in the polypeptides of the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, Science 244: 1081-1085, 1989; Bass et al., Proc. Natl. Acad. Sci. USA 88:4498-4502, 1991). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity as disclosed below to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., J. Biol. Chem. 271:4699-4708, 1996. Sites of ligand-receptor interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., Science 255:306-312, 1992; Smith et al., J. Mol. Biol. 224:899-904, 1992; Wlodaver et al., FEBS Lett. 309:59-64, 1992.

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis and screening, such as those disclosed by Reidhaar-Olson and Sauer (Science 241:53-57, 1988) or Bowie and Sauer (Proc. Natl. Acad. Sci. USA 86:2152-2156, 1989). Briefly, these authors disclose methods for simultaneously randomizing two or more positions in a polypeptide, selecting for functional polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman et al., Biochem. 30:10832-10837, 1991; Ladner et al., U.S.

Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204) and region-directed mutagenesis (Derbyshire et al., Gene 46:145, 1986; Ner et al., DNA 7:127, 1988).

Amino acid sequence changes are made in zsig15 polypeptides so as to minimize disruption of higher order structure essential to biological activity. In this regard, it is generally preferred to retain the overall hydrophilicity profile of the natural sequence. A hydrophilicity profile of the sequence shown in SEQ ID NO:2 is shown in Fig. 1.

Variants of the disclosed zsig15 DNA and polypeptide sequences can be generated through DNA shuffling as disclosed by Stemmer, Nature 370:389-91, 1994, Stemmer, Proc. Natl. Acad. Sci. USA 91:10747-51, 1994 and WIPO Publication WO 97/20078. Briefly, variant DNAs are generated by *in vitro* homologous recombination by random fragmentation of a parent DNA followed by reassembly using PCR, resulting in randomly introduced point mutations. This technique can be modified by using a family of parent DNAs, such as allelic variants or DNAs from different species, to introduce additional variability into the process. Selection or screening for the desired activity, followed by additional iterations of mutagenesis and assay provides for rapid "evolution" of sequences by selecting for desirable mutations while simultaneously selecting against detrimental changes.

Mutagenesis methods as disclosed herein can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides in host cells. Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using modern equipment. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

Using the methods discussed herein, one of ordinary skill in the art can identify and/or prepare a variety of polypeptide fragments or variants of SEQ ID NO:2 or that retain the properties of the wild-type zsig15 protein.

The zsig15 polypeptides of the present invention, including full-length polypeptides, biologically active fragments, and fusion polypeptides, can be produced in genetically engineered host cells according to conventional techniques. Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA and grown in culture, and include bacteria, fungal cells, and cultured higher eukaryotic cells. Eukaryotic cells, particularly cultured cells of multicellular organisms, are preferred. Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989, and Ausubel et al., eds., Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY, 1987.

In general, a DNA sequence encoding a zsig15 polypeptide is operably linked to other genetic elements required for its expression, generally including a transcription promoter and terminator, within an expression vector. The vector will also commonly contain one or more selectable markers and one or more origins of replication, although those skilled in the art will recognize that within certain systems selectable markers may be provided on separate vectors, and replication of the exogenous DNA may be provided by integration into the host cell genome. Selection of promoters, terminators, selectable markers, vectors and other elements is a matter of routine design within the level of ordinary skill in the art. Many such elements are described in

the literature and are available through commercial suppliers.

To direct a zsig15 polypeptide into the secretory pathway of a host cell, a secretory signal
5 sequence (also known as a leader sequence, prepro sequence or pre sequence) is provided in the expression vector. The secretory signal sequence may be that of zsig15, or may be derived from another secreted protein (e.g., t-PA) or synthesized *de novo*. The secretory
10 signal sequence is operably linked to the zsig15 DNA sequence, i.e., the two sequences are joined in the correct reading frame and positioned to direct the newly synthesized polypeptide into the secretory pathway of the host cell. Secretory signal sequences are commonly
15 positioned 5' to the DNA sequence encoding the polypeptide of interest, although certain secretory signal sequences may be positioned elsewhere in the DNA sequence of interest (see, e.g., Welch et al., U.S. Patent No. 5,037,743; Holland et al., U.S. Patent No.
20 5,143,830).

Alternatively, the secretory signal sequence contained in the polypeptides of the present invention is used to direct other polypeptides into the secretory pathway. The present invention provides for such fusion
25 polypeptides. A signal fusion polypeptide can be made wherein a secretory signal sequence derived from amino acid residues 1 to 15 through 21 of SEQ ID NO:2 is be operably linked to another polypeptide using methods known in the art and disclosed herein. The secretory
30 signal sequence contained in the fusion polypeptides of the present invention is preferably fused amino-terminally to an additional peptide to direct the additional peptide into the secretory pathway. Such constructs have numerous applications known in the art.
35 For example, these novel secretory signal sequence fusion constructs can direct the secretion of an active

component of a normally non-secreted protein, such as a receptor. Such fusions may be used *in vivo* or *in vitro* to direct peptides through the secretory pathway.

Cultured mammalian cells are suitable hosts
5 within the present invention. Methods for introducing exogenous DNA into mammalian host cells include calcium phosphate-mediated transfection (Wigler et al., Cell 14:725, 1978; Corsaro and Pearson, Somatic Cell Genetics 7:603, 1981; Graham and Van der Eb, Virology 52:456,
10 1973), electroporation (Neumann et al., EMBO J. 1:841-845, 1982), DEAE-dextran mediated transfection (Ausubel et al., ibid.), and liposome-mediated transfection (Hawley-Nelson et al., Focus 15:73, 1993; Ciccarone et al., Focus 15:80, 1993), and viral vectors (Miller and
15 Rosman, BioTechniques 7:980-90, 1989; Wang and Finer, Nature Med. 2:714-716, 1996). The production of recombinant polypeptides in cultured mammalian cells is disclosed, for example, by Levinson et al., U.S. Patent No. 4,713,339; Hagen et al., U.S. Patent No. 4,784,950;
20 Palmiter et al., U.S. Patent No. 4,579,821; and Ringold, U.S. Patent No. 4,656,134. Suitable cultured mammalian cells include the COS-1 (ATCC No. CRL 1650), COS-7 (ATCC No. CRL 1651), BHK (ATCC No. CRL 1632), BHK 570 (ATCC No. CRL 10314), 293 (ATCC No. CRL 1573; Graham et al., J.
25 Gen. Virol. 36:59-72, 1977) and Chinese hamster ovary (e.g. CHO-K1; ATCC No. CCL 61) cell lines. Additional suitable cell lines are known in the art and available from public depositories such as the American Type Culture Collection, Rockville, Maryland. In general,
30 strong transcription promoters are preferred, such as promoters from SV-40 or cytomegalovirus. See, e.g., U.S. Patent No. 4,956,288. Other suitable promoters include those from metallothionein genes (U.S. Patent Nos. 4,579,821 and 4,601,978) and the adenovirus major late
35 promoter.

Drug selection is generally used to select for cultured mammalian cells into which foreign DNA has been inserted. Such cells are commonly referred to as "transfectants". Cells that have been cultured in the presence of the selective agent and are able to pass the gene of interest to their progeny are referred to as "stable transfectants." A preferred selectable marker is a gene encoding resistance to the antibiotic neomycin. Selection is carried out in the presence of a neomycin-type drug, such as G-418 or the like. Selection systems can also be used to increase the expression level of the gene of interest, a process referred to as "amplification." Amplification is carried out by culturing transfectants in the presence of a low level of the selective agent and then increasing the amount of selective agent to select for cells that produce high levels of the products of the introduced genes. A preferred amplifiable selectable marker is dihydrofolate reductase, which confers resistance to methotrexate. Other drug resistance genes (e.g., hygromycin resistance, multi-drug resistance, puromycin acetyltransferase) can also be used. Alternative markers that introduce an altered phenotype, such as green fluorescent protein, or cell surface proteins such as CD4, CD8, Class I MHC, placental alkaline phosphatase may be used to sort transfected cells from untransfected cells by such means as FACS sorting or magnetic bead separation technology.

Other higher eukaryotic cells can also be used as hosts, including plant cells, insect cells and avian cells. The use of *Agrobacterium rhizogenes* as a vector for expressing genes in plant cells has been reviewed by Sinkar et al., J. Biosci. (Bangalore) 11:47-58, 1987. Transformation of insect cells and production of foreign polypeptides therein is disclosed by Guarino et al., U.S. Patent No. 5,162,222 and WIPO publication WO 94/06463. Insect cells can be infected with recombinant

5 baculovirus, commonly derived from *Autographa californica* nuclear polyhedrosis virus (AcNPV). DNA encoding the zsig15 polypeptide is inserted into the baculoviral genome in place of the AcNPV polyhedrin gene coding sequence by one of two methods. The first is the traditional method of homologous DNA recombination between wild-type AcNPV and a transfer vector containing the zsig15 flanked by AcNPV sequences. Suitable insect cells, e.g. SF9 cells, are infected with wild-type AcNPV and transfected with a transfer vector comprising a zsig15 polynucleotide operably linked to an AcNPV polyhedrin gene promoter, terminator, and flanking sequences. See, King, L.A. and Possee, R.D., The Baculovirus Expression System: A Laboratory Guide, 15 London, Chapman & Hall; O'Reilly, D.R. et al., Baculovirus Expression Vectors: A Laboratory Manual, New York, Oxford University Press., 1994; and, Richardson, C. D., Ed., Baculovirus Expression Protocols. Methods in Molecular Biology, Totowa, NJ, Humana Press, 1995. 20 Natural recombination within an insect cell will result in a recombinant baculovirus which contains zsig15 driven by the polyhedrin promoter. Recombinant viral stocks are made by methods commonly used in the art.

The second method of making recombinant 25 baculovirus utilizes a transposon-based system described by Luckow (Luckow, V.A, et al., J Virol 67:4566-79, 1993). This system is sold in the Bac-to-Bac kit (Life Technologies, Rockville, MD). This system utilizes a transfer vector, pFastBac1™ (Life Technologies) 30 containing a Tn7 transposon to move the DNA encoding the zsig15 polypeptide into a baculovirus genome maintained in E. coli as a large plasmid called a "bacmid." The pFastBac1™ transfer vector utilizes the AcNPV polyhedrin promoter to drive the expression of the gene of interest, 35 in this case zsig15. However, pFastBac1™ can be modified to a considerable degree. The polyhedrin promoter can be

removed and substituted with the baculovirus basic protein promoter (also known as Pcor, p6.9 or MP promoter) which is expressed earlier in the baculovirus infection, and has been shown to be advantageous for
5 expressing secreted proteins. See, Hill-Perkins, M.S. and Possee, R.D., J Gen Virol 71:971-6, 1990; Bonning, B.C. et al., J Gen Virol 75:1551-6, 1994; and, Chazenbalk, G.D., and Rapoport, B., J Biol Chem 270:1543-9, 1995. In such transfer vector constructs, a short or
10 long version of the basic protein promoter can be used. Moreover, transfer vectors can be constructed which replace the native zsig15 secretory signal sequences with secretory signal sequences derived from insect proteins. For example, a secretory signal sequence from Ecdysteroid
15 Glucosyltransferase (EGT), honey bee Melittin (Invitrogen, Carlsbad, CA), or baculovirus gp67 (PharMingen, San Diego, CA) can be used in constructs to replace the native zsig15 secretory signal sequence. In addition, transfer vectors can include an in-frame fusion
20 with DNA encoding an epitope tag at the C- or N-terminus of the expressed zsig15 polypeptide, for example, a Glu-Glu epitope tag (Grussenmeyer, T. et al., Proc Natl Acad Sci. 82:7952-4, 1985). Using a technique known in the art, a transfer vector containing zsig15 is transformed
25 into E. Coli, and screened for bacmids which contain an interrupted lacZ gene indicative of recombinant baculovirus. The bacmid DNA containing the recombinant baculovirus genome is isolated, using common techniques, and used to transfect *Spodoptera frugiperda* cells, e.g.
30 Sf9 cells. Recombinant virus that expresses zsig15 is subsequently produced. Recombinant viral stocks are made by methods commonly used the art.

The recombinant virus is used to infect host cells, typically a cell line derived from the fall
35 armyworm, *Spodoptera frugiperda*. See, in general, Glick and Pasternak, Molecular Biotechnology: Principles and

Applications of Recombinant DNA, ASM Press, Washington, D.C., 1994. Another suitable cell line is the High Five⁰™ cell line (Invitrogen) derived from *Trichoplusia ni* (U.S. Patent #5,300,435). Commercially available

5 serum-free media are used to grow and maintain the cells. Suitable media are Sf900 II™ (Life Technologies) or ESF 921™ (Expression Systems) for the Sf9 cells; and Excell0405™ (JRH Biosciences, Lenexa, KS) or Express Five⁰™ (Life Technologies) for the *T. ni* cells. The

10 cells are grown up from an inoculation density of approximately $2-5 \times 10^5$ cells to a density of $1-2 \times 10^6$ cells at which time a recombinant viral stock is added at a multiplicity of infection (MOI) of 0.1 to 10, more typically near 3. The recombinant virus-infected cells

15 typically produce the recombinant zsig15 polypeptide at 12-72 hours post-infection and secrete it with varying efficiency into the medium. The culture is usually harvested 48 hours post-infection. Centrifugation is used to separate the cells from the medium (supernatant).

20 The supernatant containing the zsig15 polypeptide is filtered through micropore filters, usually 0.45 μm pore size. Procedures used are generally described in available laboratory manuals (King, L. A. and Possee, R.D., ibid.; O'Reilly, D.R. et al., ibid.; Richardson, C.

25 D., ibid.). Subsequent purification of the zsig15 polypeptide from the supernatant can be achieved using methods described herein.

Fungal cells, including yeast cells, can also be used within the present invention. Yeast species of

30 particular interest in this regard include *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Pichia methanolica*. Methods for transforming *S. cerevisiae* cells with exogenous DNA and producing recombinant polypeptides therefrom are disclosed by, for example, Kawasaki, U.S.

Patent No. 4,599,311; Kawasaki et al., U.S. Patent No. 4,931,373; Brake, U.S. Patent No. 4,870,008; Welch et al., U.S. Patent No. 5,037,743; and Murray et al., U.S. Patent No. 4,845,075. Transformed cells are selected by
5 phenotype determined by the selectable marker, commonly drug resistance or the ability to grow in the absence of a particular nutrient (e.g., leucine). A preferred vector system for use in *Saccharomyces cerevisiae* is the *POT1* vector system disclosed by Kawasaki et al. (U.S.
10 Patent No. 4,931,373), which allows transformed cells to be selected by growth in glucose-containing media. Suitable promoters and terminators for use in yeast include those from glycolytic enzyme genes (see, e.g., Kawasaki, U.S. Patent No. 4,599,311; Kingsman et al.,
15 U.S. Patent No. 4,615,974; and Bitter, U.S. Patent No. 4,977,092) and alcohol dehydrogenase genes. See also U.S. Patents Nos. 4,990,446; 5,063,154; 5,139,936 and 4,661,454. Transformation systems for other yeasts, including *Hansenula polymorpha*, *Schizosaccharomyces*
20 *pombe*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, *Ustilago maydis*, *Pichia pastoris*, *Pichia methanolica*, *Pichia guilliermondii* and *Candida maltosa* are known in the art. See, for example, Gleeson et al., J. Gen. Microbiol. 132:3459-3465, 1986 and Cregg, U.S. Patent No.
25 4,882,279. *Aspergillus* cells may be utilized according to the methods of McKnight et al., U.S. Patent No. 4,935,349. Methods for transforming *Acremonium chrysogenum* are disclosed by Sumino et al., U.S. Patent No. 5,162,228. Methods for transforming *Neurospora* are
30 disclosed by Lambowitz, U.S. Patent No. 4,486,533.

The use of *Pichia methanolica* as host for the production of recombinant proteins is disclosed in WIPO Publications WO 97/17450, WO 97/17451, WO 98/02536, and WO 98/02565. DNA molecules for use in transforming *P.*

methanolica will commonly be prepared as double-stranded, circular plasmids, which are preferably linearized prior to transformation. For polypeptide production in *P. methanolica*, it is preferred that the promoter and terminator in the plasmid be that of a *P. methanolica* gene, such as a *P. methanolica* alcohol utilization gene (*AUG1* or *AUG2*). Other useful promoters include those of the dihydroxyacetone synthase (*DHAS*), formate dehydrogenase (*FMD*), and catalase (*CAT*) genes. To facilitate integration of the DNA into the host chromosome, it is preferred to have the entire expression segment of the plasmid flanked at both ends by host DNA sequences. A preferred selectable marker for use in *Pichia methanolica* is a *P. methanolica ADE2* gene, which encodes phosphoribosyl-5-aminoimidazole carboxylase (*AIRC*; EC 4.1.1.21), which allows *ade2* host cells to grow in the absence of adenine. For large-scale, industrial processes where it is desirable to minimize the use of methanol, it is preferred to use host cells in which both methanol utilization genes (*AUG1* and *AUG2*) are deleted. For production of secreted proteins, host cells deficient in vacuolar protease genes (*PEP4* and *PRB1*) are preferred. Electroporation is used to facilitate the introduction of a plasmid containing DNA encoding a polypeptide of interest into *P. methanolica* cells. It is preferred to transform *P. methanolica* cells by electroporation using an exponentially decaying, pulsed electric field having a field strength of from 2.5 to 4.5 kV/cm, preferably about 3.75 kV/cm, and a time constant (τ) of from 1 to 40 milliseconds, most preferably about 20 milliseconds.

Prokaryotic host cells, including strains of the bacteria *Escherichia coli*, *Bacillus* and other genera are also useful host cells within the present invention. Techniques for transforming these hosts and expressing

foreign DNA sequences cloned therein are well known in the art (see, e.g., Sambrook et al., ibid.). When expressing a zsig15 polypeptide in bacteria such as *E. coli*, the polypeptide may be retained in the cytoplasm, typically as insoluble granules, or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed, and the granules are recovered and denatured using, for example, guanidine isothiocyanate or urea. The denatured polypeptide can then be refolded and dimerized by diluting the denaturant, such as by dialysis against a solution of urea and a combination of reduced and oxidized glutathione, followed by dialysis against a buffered saline solution. In the latter case, the polypeptide can be recovered from the periplasmic space in a soluble and functional form by disrupting the cells (by, for example, sonication or osmotic shock) to release the contents of the periplasmic space and recovering the protein, thereby obviating the need for denaturation and refolding.

Transformed or transfected host cells are cultured according to conventional procedures in a culture medium containing nutrients and other components required for the growth of the chosen host cells. A variety of suitable media, including defined media and complex media, are known in the art and generally include a carbon source, a nitrogen source, essential amino acids, vitamins and minerals. Media may also contain such components as growth factors or serum, as required. The growth medium will generally select for cells containing the exogenously added DNA by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker carried on the expression vector or co-transfected into the host cell.

P. methanolica cells are cultured in a medium comprising adequate sources of carbon, nitrogen, and

trace nutrients at a temperature of about 25°C to 35°C. Liquid cultures are provided with sufficient aeration by conventional means, such as shaking of small flasks or sparging of fermentors. A preferred culture medium for
5 *P. methanolica* is YEPD (2% D-glucose, 2% Bacto™ Peptone (Difco Laboratories, Detroit, MI), 1% Bacto™ yeast extract (Difco Laboratories), 0.004% adenine and 0.006% L-leucine).

zsig15 polypeptides or fragments thereof may
10 also be prepared through chemical synthesis. zsig15 polypeptides may be monomers or multimers; glycosylated or non-glycosylated; pegylated or non-pegylated; and may or may not include an initial methionine amino acid residue.

15 Polypeptides of the present invention are useful for measuring changes in levels of expression zsig15 polypeptides. Because zsig15 expression is restricted to specific tissues (i.e., prostate, colon and breast) and bone marrow, changes in expression levels
20 could be used to monitor metabolism within these tissues. For example, increases in expression and/or transcription of zsig15 polypeptides and polynucleotides, may be predictive for increased cell proliferation of tumor cells. Furthermore, expression of zsig15 in tissue not
25 normally expressing zsig15, may be indicative of metastases of tumor cells.

Zsig15 has been demonstrated to be expressed differentially in certain epithelial tissues and carcinomas, particularly in colon and breast.
30 Differential expression is the transient expression, or lack thereof, of specific genes, proteins or other phenotypic properties (known as differentiation markers) that occur during the progress of maturation in a cell or tissue. A set of differentiation markers is defined as
35 one or more phenotypic properties that can be identified and are specific to a particular cell type. Thus,

pluripotent stem cells that can regenerate without commitment to a lineage express a set of differentiation markers that are lost when commitment to a cell lineage is made. Precursor cells express a set of
5 differentiation markers that may or may not continue to be expressed as the cells progress down the cell lineage pathway toward maturation. Differentiation markers that are expressed exclusively by mature cells are usually functional properties such as cell products, enzymes to
10 produce cell products and receptors.

Zsig15 expression can be used as a differentiation marker in normal and tumor tissues to determine the stage of the tumor or maturity of a cell. Zsig15 will be particularly valuable as a marker for
15 epithelial cells and tumor of epithelial origin, and more particularly epithelial cells and epithelial-derived tumors from colon, breast or prostate tissues.

A set of differentiation markers is defined as one or more phenotypic properties that can be identified
20 and are specific to a particular cell type. Differentiation markers are transiently exhibited at various stages of cell lineage. Pluripotent stem cells that can regenerate without commitment to a lineage express a set of differentiation markers that are lost
25 when commitment to a cell lineage is made. Precursor cells express a set of differentiation markers that may or may not continue to be expressed as the cells progress down the cell lineage pathway toward maturation. Differentiation markers that are expressed exclusively by
30 mature cells are usually functional properties such as cell products, enzymes to produce cell products and receptors.

The activity of molecules of the present invention can be measured using a variety of assays that
35 measure proliferation and/or differentiation of specific cell types, chemotaxis, adhesion, changes in ion channel

influx, regulation of second messenger levels and neurotransmitter release. Such assays are well known in the art. See, for example, in "Basic & Clinical Endocrinology Ser., Vol. Vol. 3," Cytochemical Bioassays: Techniques & Applications, Chayen; Chayen, Bitensky, eds., Dekker, New York, 1983.

Proliferation and differentiation can be measured using cultured cells or *in vivo* by administering molecules of the claimed invention to the appropriate animal model. Assays measuring cell proliferation or differentiation are well known in the art. For example, assays measuring proliferation include such assays as chemosensitivity to neutral red dye (Cavanaugh et al., Investigational New Drugs 8:347-354, 1990, incorporated herein by reference), incorporation of radiolabelled nucleotides (Cook et al., Analytical Biochem. 179:1-7, 1989, incorporated herein by reference), incorporation of 5-bromo-2'-deoxyuridine (BrdU) in the DNA of proliferating cells (Porstmann et al., J. Immunol. Methods 82:169-179, 1985, incorporated herein by reference), and use of tetrazolium salts (Mosmann, J. Immunol. Methods 65:55-63, 1983; Alley et al., Cancer Res. 48:589-601, 1988; Marshall et al., Growth Reg. 5:69-84, 1995; and Scudiero et al., Cancer Res. 48:4827-4833, 1988; all incorporated herein by reference). Assays measuring differentiation include, for example, measuring cell-surface markers associated with stage-specific expression of a tissue, enzymatic activity, functional activity or morphological changes (Watt, FASEB, 5:281-284, 1991; Francis, Differentiation 57:63-75, 1994; Raes, Adv. Anim. Cell Biol. Technol. Bioprocesses, 161-171, 1989; all incorporated herein by reference).

Polypeptides of the present invention can be measured *in vitro* using cultured cells or *in vivo* by administering molecules of the claimed invention to the appropriate animal model. For instance, BHK transfected

expression host cells may be embedded in an alginate environment and injected (implanted) into recipient animals. Alginate-poly-L-lysine microencapsulation, permselective membrane encapsulation and diffusion chambers have been described as a means to entrap transfected mammalian cells or primary mammalian cells. These types of non-immunogenic "encapsulations" or microenvironments permit the transfer of nutrients into the microenvironment, and also permit the diffusion of proteins and other macromolecules secreted or released by the captured cells across the environmental barrier to the recipient animal. Most importantly, the capsules or microenvironments mask and shield the foreign, embedded cells from the recipient animal's immune response. Such microenvironments can extend the life of the injected cells from a few hours or days (naked cells) to several weeks (embedded cells).

Alginate threads provide a simple and quick means for generating embedded cells. The materials needed to generate the alginate threads are readily available and relatively inexpensive. Once made, the alginate threads are relatively strong and durable, both *in vitro* and, based on data obtained using the threads, *in vivo*. The alginate threads are easily manipulable and the methodology is scalable for preparation of numerous threads. In an exemplary procedure, 3% alginate is prepared in sterile H₂O, and sterile filtered. Just prior to preparation of alginate threads, the alginate solution is again filtered. An approximately 50% cell suspension (containing about 5×10^5 to about 5×10^7 cells/ml) is mixed with the 3% alginate solution. One ml of the alginate/cell suspension is extruded into a 100 mM sterile filtered CaCl₂ solution over a time period of ~15 min, forming a "thread". The extruded thread is then transferred into a solution of 50 mM CaCl₂, and then into a solution of 25 mM CaCl₂. The thread is then rinsed

with deionized water before coating the thread by incubating in a 0.01% solution of poly-L-lysine. Finally, the thread is rinsed with Lactated Ringer's Solution and drawn from solution into a syringe barrel
5 (without needle attached). A large bore needle is then attached to the syringe, and the thread is intraperitoneally injected into a recipient in a minimal volume of the Lactated Ringer's Solution.

An alternative *in vivo* approach for assaying
10 proteins of the present invention involves viral delivery systems. Exemplary viruses for this purpose include adenovirus, herpesvirus, vaccinia virus and adeno-associated virus (AAV). Adenovirus, a double-stranded DNA virus, is currently the best studied gene transfer
15 vector for delivery of heterologous nucleic acid (for a review, see T.C. Becker et al., Meth. Cell Biol. 43:161-89, 1994; and J.T. Douglas and D.T. Curiel, Science & Medicine 4:44-53, 1997). The adenovirus system offers several advantages: adenovirus can (i) accommodate
20 relatively large DNA inserts; (ii) be grown to high-titer; (iii) infect a broad range of mammalian cell types; and (iv) be used with a large number of available vectors containing different promoters. Also, because adenoviruses are stable in the bloodstream, they can be
25 administered by intravenous injection.

By deleting portions of the adenovirus genome, larger inserts (up to 7 kb) of heterologous DNA can be accommodated. These inserts can be incorporated into the viral DNA by direct ligation or by homologous
30 recombination with a co-transfected plasmid. In an exemplary system, the essential E1 gene has been deleted from the viral vector, and the virus will not replicate unless the E1 gene is provided by the host cell (the human 293 cell line is exemplary). When intravenously
35 administered to intact animals, adenovirus primarily targets the liver. If the adenoviral delivery system has

an E1 gene deletion, the virus cannot replicate in the host cells. However, the host's tissue (e.g., liver) will express and process (and, if a secretory signal sequence is present, secrete) the heterologous protein.

- 5 Secreted proteins will enter the circulation in the highly vascularized liver, and effects on the infected animal can be determined.

The adenovirus system can also be used for protein production *in vitro*. By culturing adenovirus-
10 infected non-293 cells under conditions where the cells are not rapidly dividing, the cells can produce proteins for extended periods of time. For instance, BHK cells are grown to confluence in cell factories, then exposed to the adenoviral vector encoding the secreted protein of
15 interest. The cells are then grown under serum-free conditions, which allows infected cells to survive for several weeks without significant cell division. Alternatively, adenovirus vector infected 293S cells can be grown in suspension culture at relatively high cell
20 density to produce significant amounts of protein (see Garnier et al., Cytotechnol. 15:145-55, 1994). With either protocol, an expressed, secreted heterologous protein can be repeatedly isolated from the cell culture supernatant. Within the infected 293S cell production
25 protocol, non-secreted proteins may also be effectively obtained.

Expressed recombinant zsig15 polypeptides (or chimeric zsig15 polypeptides) can be purified using fractionation and/or conventional purification methods
30 and media. Ammonium sulfate precipitation and acid or chaotrope extraction may be used for fractionation of samples. Exemplary purification steps may include hydroxyapatite, size exclusion, FPLC and reverse-phase high performance liquid chromatography. Suitable
35 chromatographic media include derivatized dextrans, agarose, cellulose, polyacrylamide, specialty silicas,

and the like. PEI, DEAE, QAE and Q derivatives are preferred. Exemplary chromatographic media include those media derivatized with phenyl, butyl, or octyl groups, such as Phenyl-Sepharose FF (Pharmacia), Toyopearl butyl 5 650 (Toso Haas, Montgomeryville, PA), Octyl-Sepharose (Pharmacia) and the like; or polyacrylic resins, such as Amberchrom CG 71 (Toso Haas) and the like. Suitable solid supports include glass beads, silica-based resins, cellulosic resins, agarose beads, cross-linked agarose 10 beads, polystyrene beads, cross-linked polyacrylamide resins and the like that are insoluble under the conditions in which they are to be used. These supports may be modified with reactive groups that allow attachment of proteins by amino groups, carboxyl groups, 15 sulfhydryl groups, hydroxyl groups and/or carbohydrate moieties. Examples of coupling chemistries include cyanogen bromide activation, N-hydroxysuccinimide activation, epoxide activation, sulfhydryl activation, hydrazide activation, and carboxyl and amino derivatives 20 for carbodiimide coupling chemistries. These and other solid media are well known and widely used in the art, and are available from commercial suppliers. Methods for binding receptor polypeptides to support media are well known in the art. Selection of a particular method is a 25 matter of routine design and is determined in part by the properties of the chosen support. See, for example, Affinity Chromatography: Principles & Methods, Pharmacia LKB Biotechnology, Uppsala, Sweden, 1988.

The polypeptides of the present invention can 30 be isolated by exploitation of specific properties. For example, immobilized metal ion adsorption (IMAC) chromatography can be used to purify histidine-rich proteins, including those comprising polyhistidine tags. Briefly, a gel is first charged with divalent metal ions 35 to form a chelate (Sulkowski, Trends in Biochem. 3:1-7, 1985). Histidine-rich proteins will be adsorbed to this

matrix with differing affinities, depending upon the metal ion used, and will be eluted by competitive elution, lowering the pH, or use of strong chelating agents. Other methods of purification include
5 purification of glycosylated proteins by lectin affinity chromatography and ion exchange chromatography (Methods in Enzymol., Vol. 182, "Guide to Protein Purification", M. Deutscher, (ed.), Acad. Press, San Diego, 1990, pp.529-39). Within additional embodiments of the
10 invention, a fusion of the polypeptide of interest and an affinity tag (e.g., maltose-binding protein, an immunoglobulin domain) may be constructed to facilitate purification.

Using methods described in the art, polypeptide
15 fusions, or hybrid zsig15 proteins, are constructed using regions or domains of the inventive zsig15 in combination with those of other heterologous proteins (Sambrook et al., ibid., Altschul et al., ibid., Picard, Cur. Opin. Biology, 5:511-5, 1994, and references therein). These
20 methods allow the determination of the biological importance of larger domains or regions in a polypeptide of interest. Such hybrids may alter reaction kinetics, binding, constrict or expand the substrate specificity, or alter tissue and cellular localization of a
25 polypeptide, and can be applied to polypeptides of unknown structure.

Fusion proteins can be prepared by methods known to those skilled in the art by preparing each component of the fusion protein and chemically
30 conjugating them. Alternatively, a polynucleotide encoding both components of the fusion protein in the proper reading frame can be generated using known techniques and expressed by the methods described herein. For example, part or all of a domain(s) conferring a
35 biological function may be swapped between zsig15 of the present invention with the functionally equivalent

domain(s) from another family member. Such domains include, but are not limited to, the secretory signal sequence and conserved motifs. Such fusion proteins would be expected to have a biological functional profile that is the same or similar to polypeptides of the present invention or other family, depending on the fusion constructed. Moreover, such fusion proteins may exhibit other properties as disclosed herein.

The gene for molecules of the present invention has been mapped to the chromosomal location 19Q 13.1-13.2. This chromosomal location is distinguished by a clustering of multiple genes, that when expressed at high level, are associated with carcinomas. Kallikrein/Prostate-specific antigen (PSA) is localized to chromosome 19Q 13.12-13.4 and has been shown to be overexpressed in prostate cancers (Winderickx et al., Mol. Cell. Endocrinol. 62:217-226, 1989). PSA is a single chain glycoprotein of 33 kDa. Another cluster of genes localized to chromosome 19Q 13.1-13.3 is the pregnancy-specific beta glycoprotein family (PSBGs). PSBGs have been used to diagnose pregnancy, hydatidiform moles and choriocarcinoma (McLenachan et al., Genomics 22:356-363, 1994). The PSBGs are members of the immunoglobulin super family and related to another gene family found in this cluster, carcinoembryonic antigen (CEA). CEA gene family is located on the long arm of chromosome 19 and has been identified as a marker for colorectal, gastric and pancreatic carcinomas (Hodge, Cancer Immunol. Immunother. 43:127-143, 1996). Thus, the gene of the present invention provides an additional tool for further elucidation of a chromosomal location that is believed to be important in diagnosis of various cancers.

In addition, family studies can be done to identify whether the gene for zsig15 is mutated and therefore indicative of an inherited form of disease. Mutations of genes can identified by methods well known

to those with ordinary skill in such arts. For example, nucleic acids (i.e., genomic DNA, cDNA or mRNA) are extracted from tissue samples or biological fluids and analyzed by PCR using PCR primers complementary to
5 nucleic acids of the present invention. Analyses involve identification of insertions, deletions and substitutions that may, for example, differ in mobility when subjected to gel electrophoresis or can be cleaved by ribonuclease at single base mismatch in a RNA-DNA heteroduplex (Myers
10 et al., Science, 230:1242-1246, 1985).

zsig15 polypeptides can also be used to prepare antibodies that specifically bind to zsig15 epitopes, peptides or polypeptides. Methods for preparing polyclonal and monoclonal antibodies are well known in
15 the art.

In view of the tissue distribution observed for zsig15, agonists (including the natural ligand/substrate/ cofactor/ etc.) and antagonists have potential in both *in vitro* and *in vivo* applications. For example,
20 zsig15 and agonist compounds are useful as components of defined cell culture media, and may be used alone or in combination with other cytokines and hormones to replace serum that is commonly used in cell culture. Agonists are thus useful in specifically promoting the growth
25 and/or development of epithelial cells, particularly of colon or breast tissue origin.

Antagonists are also useful as research reagents for characterizing sites of ligand-receptor interaction. Inhibitors of zsig15 activity (zsig15
30 antagonists) include anti-zsig15 antibodies and soluble zsig15 receptors, as well as other peptidic and non-peptidic agents (including ribozymes).

zsig15 can also be used to identify inhibitors (antagonists) of its activity. Test compounds are added
35 to the assays disclosed herein to identify compounds that inhibit the activity of zsig15. In addition to those

assays disclosed herein, samples can be tested for inhibition of zsig15 activity within a variety of assays designed to measure receptor binding or the stimulation/inhibition of zsig15-dependent cellular responses. For example, zsig15-responsive cell lines can be transfected with a reporter gene construct that is responsive to a zsig15-stimulated cellular pathway. Reporter gene constructs of this type are known in the art, and will generally comprise a zsig15-DNA response element operably linked to a gene encoding an assayable protein, such as luciferase. DNA response elements can include, but are not limited to, cyclic AMP response elements (CRE), hormone response elements (HRE) insulin response element (IRE) (Nasrin et al., Proc. Natl. Acad. Sci. USA 87:5273-7, 1990) and serum response elements (SRE) (Shaw et al. Cell 56: 563-72, 1989). Cyclic AMP response elements are reviewed in Roestler et al., J. Biol. Chem. 263 (19):9063-6; 1988 and Habener, Molec. Endocrinol. 4 (8):1087-94; 1990. Hormone response elements are reviewed in Beato, Cell 56:335-44; 1989. Candidate compounds, solutions, mixtures or extracts are tested for the ability to inhibit the activity of zsig15 on the target cells as evidenced by a decrease in zsig15 stimulation of reporter gene expression. Assays of this type will detect compounds that directly block zsig15 binding to cell-surface receptors, as well as compounds that block processes in the cellular pathway subsequent to receptor-ligand binding. In the alternative, compounds or other samples can be tested for direct blocking of zsig15 binding to receptor using zsig15 tagged with a detectable label (e.g., ¹²⁵I, biotin, horseradish peroxidase, FITC, and the like). Within assays of this type, the ability of a test sample to inhibit the binding of labeled zsig15 to the receptor is indicative of inhibitory activity, which can be confirmed through secondary assays. Receptors used within binding

assays may be cellular receptors or isolated, immobilized receptors.

A zsig15 polypeptide, or portions thereof, can be expressed as a fusion with an immunoglobulin heavy chain constant region, typically an F_C fragment, which contains two constant region domains and lacks the variable region. Methods for preparing such fusions are disclosed in U.S. Patents Nos. 5,155,027 and 5,567,584. Such fusions are typically secreted as multimeric molecules wherein the F_C portions are disulfide bonded to each other and two non-Ig polypeptides are arrayed in closed proximity to each other. Fusions of this type can be used to affinity purify ligand, as *in vitro* assay tool, or antagonist. For use in assays, the chimeras are bound to a support via the F_C region and used in an ELISA format.

A zsig15 ligand-binding polypeptide can also be used for purification of ligand. The polypeptide is immobilized on a solid support, such as beads of agarose, cross-linked agarose, glass, cellulosic resins, silica-based resins, polystyrene, cross-linked polyacrylamide, or like materials that are stable under the conditions of use. Methods for linking polypeptides to solid supports are known in the art, and include amine chemistry, cyanogen bromide activation, N-hydroxysuccinimide activation, epoxide activation, sulfhydryl activation, and hydrazide activation. The resulting medium will generally be configured in the form of a column, and fluids containing ligand are passed through the column one or more times to allow ligand to bind to the receptor polypeptide. The ligand is then eluted using changes in salt concentration, chaotropic agents (guanidine HCl), or pH to disrupt ligand-receptor binding.

An assay system that uses a ligand-binding receptor (or an antibody, one member of a complement/anti-complement pair) or a binding fragment thereof, and

a commercially available biosensor instrument (BIAcore, Pharmacia Biosensor, Piscataway, NJ) may be advantageously employed. Such receptor, antibody, member of a complement/anti-complement pair or fragment is immobilized onto the surface of a receptor chip. Use of this instrument is disclosed by Karlsson, J. Immunol. Methods 145:229-40, 1991 and Cunningham and Wells, J. Mol. Biol. 234:554-63, 1993. A receptor, antibody, member or fragment is covalently attached, using amine or sulfhydryl chemistry, to dextran fibers that are attached to gold film within the flow cell. A test sample is passed through the cell. If a ligand, epitope, or opposite member of the complement/anti-complement pair is present in the sample, it will bind to the immobilized receptor, antibody or member, respectively, causing a change in the refractive index of the medium, which is detected as a change in surface plasmon resonance of the gold film. This system allows the determination of on- and off-rates, from which binding affinity can be calculated, and assessment of stoichiometry of binding.

Ligand-binding receptor polypeptides can also be used within other assay systems known in the art. Such systems include Scatchard analysis for determination of binding affinity (see Scatchard, Ann. NY Acad. Sci. 51: 660-72, 1949) and calorimetric assays (Cunningham et al., Science 253:545-48, 1991; Cunningham et al., Science 245:821-25, 1991).

zsig15 polypeptides can also be used to prepare antibodies that specifically bind to zsig15 epitopes, peptides or polypeptides. The zsig15 polypeptide or a fragment thereof serves as an antigen (immunogen) to inoculate an animal and elicit an immune response. Antibodies generated from this immune response can be isolated and purified as described herein. Methods for preparing and isolating polyclonal and monoclonal antibodies are well known in the art. See, for example,

Current Protocols in Immunology, Cooligan, et al. (eds.), National Institutes of Health, John Wiley and Sons, Inc., 1995; Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY, 1989; and
5 Hurrell, J. G. R., Ed., Monoclonal Hybridoma Antibodies: Techniques and Applications, CRC Press, Inc., Boca Raton, FL, 1982.

As would be evident to one of ordinary skill in the art, polyclonal antibodies can be generated from
10 inoculating a variety of warm-blooded animals, such as horses, cows, goats, sheep, dogs, chickens, rabbits, mice, and rats with a zsig15 polypeptide or a fragment thereof. The immunogenicity of a zsig15 polypeptide may be increased through the use of an adjuvant, such as alum
15 (aluminum hydroxide) or Freund's complete or incomplete adjuvant. Polypeptides useful for immunization also include fusion polypeptides, such as fusions of zsig15 or a portion thereof with an immunoglobulin polypeptide or with maltose binding protein. The polypeptide immunogen
20 may be a full-length molecule or a portion thereof. If the polypeptide portion is "haptten-like", such portion may be advantageously joined or linked to a macromolecular carrier (such as keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA) or tetanus toxoid) for
25 immunization.

As used herein, the term "antibodies" includes polyclonal antibodies, affinity-purified polyclonal antibodies, monoclonal antibodies, and antigen-binding fragments, such as F(ab')₂ and Fab proteolytic fragments.
30 Genetically engineered intact antibodies or fragments, such as chimeric antibodies, Fv fragments, single chain antibodies and the like, as well as synthetic antigen-binding peptides and polypeptides, are also included. Non-human antibodies may be humanized by grafting non-

human CDRs onto human framework and constant regions, or by incorporating the entire non-human variable domains (optionally "cloaking" them with a human-like surface by replacement of exposed residues, wherein the result is a "veneered" antibody). In some instances, humanized antibodies may retain non-human residues within the human variable region framework domains to enhance proper binding characteristics. Through humanizing antibodies, biological half-life may be increased, and the potential for adverse immune reactions upon administration to humans is reduced.

Alternative techniques for generating or selecting antibodies useful herein include *in vitro* exposure of lymphocytes to zsig15 protein or peptide, and selection of antibody display libraries in phage or similar vectors (for instance, through use of immobilized or labeled zsig15 protein or peptide). Genes encoding polypeptides having potential zsig15 polypeptide binding domains can be obtained by screening random peptide libraries displayed on phage (phage display) or on bacteria, such as *E. coli*. Nucleotide sequences encoding the polypeptides can be obtained in a number of ways, such as through random mutagenesis and random polynucleotide synthesis. These random peptide display libraries can be used to screen for peptides which interact with a known target which can be a protein or polypeptide, such as a ligand or receptor, a biological or synthetic macromolecule, or organic or inorganic substances. Techniques for creating and screening such random peptide display libraries are known in the art (Ladner et al., US Patent NO. 5,223,409; Ladner et al., US Patent NO. 4,946,778; Ladner et al., US Patent NO. 5,403,484 and Ladner et al., US Patent NO. 5,571,698) and random peptide display libraries and kits for screening such libraries are available commercially, for instance

from Clontech (Palo Alto, CA), Invitrogen Inc. (San Diego, CA), New England Biolabs, Inc. (Beverly, MA) and Pharmacia LKB Biotechnology Inc. (Piscataway, NJ). Random peptide display libraries can be screened using
5 the zsig15 sequences disclosed herein to identify proteins which bind to zsig15. These "binding proteins" which interact with zsig15 polypeptides can be used for tagging cells; for isolating homolog polypeptides by affinity purification; they can be directly or indirectly
10 conjugated to drugs, toxins, radionuclides and the like. These binding proteins can also be used in analytical methods such as for screening expression libraries and neutralizing activity. The binding proteins can also be used for diagnostic assays for determining circulating
15 levels of polypeptides; for detecting or quantitating soluble polypeptides as marker of underlying pathology or disease. These binding proteins can also act as zsig15 "antagonists" to block zsig15 binding and signal transduction *in vitro* and *in vivo*. These anti-zsig15
20 binding proteins would be useful for inhibiting ???.

Antibodies are determined to be specifically binding if: 1) they exhibit a threshold level of binding activity, and/or 2) they do not significantly cross-react with related polypeptide molecules. First, antibodies
25 herein specifically bind if they bind to a zsig15 polypeptide, peptide or epitope with a binding affinity (K_a) of 10^6 M^{-1} or greater, preferably 10^7 M^{-1} or greater, more preferably 10^8 M^{-1} or greater, and most preferably 10^9 M^{-1} or greater. The binding affinity of
30 an antibody can be readily determined by one of ordinary skill in the art, for example, by Scatchard analysis (Scatchard, G., Ann. NY Acad. Sci. 51: 660-672, 1949).

Second, antibodies are determined to specifically bind if they do not significantly cross-

react with related polypeptides. Antibodies do not significantly cross-react with related polypeptide molecules, for example, if they detect zsig15 but not known related polypeptides using a standard Western blot analysis (Ausubel et al., ibid.). Examples of known related polypeptides are orthologs, proteins from the same species that are members of a protein family (e.g. IL-16), zsig15 polypeptides, and non-human zsig15. Moreover, antibodies may be "screened against" known related polypeptides to isolate a population that specifically binds to the inventive polypeptides. For example, antibodies raised to zsig15 are adsorbed to related polypeptides adhered to insoluble matrix; antibodies specific to zsig15 will flow through the matrix under the proper buffer conditions. Such screening allows isolation of polyclonal and monoclonal antibodies non-crossreactive to closely related polypeptides (Antibodies: A Laboratory Manual, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988; Current Protocols in Immunology, Cooligan, et al. (eds.), National Institutes of Health, John Wiley and Sons, Inc., 1995). Screening and isolation of specific antibodies is well known in the art. See, Fundamental Immunology, Paul (eds.), Raven Press, 1993; Getzoff et al., Adv. in Immunol. 43: 1-98, 1988; Monoclonal Antibodies: Principles and Practice, Goding, J.W. (eds.), Academic Press Ltd., 1996; Benjamin et al., Ann. Rev. Immunol. 2: 67-101, 1984.

A variety of assays known to those skilled in the art can be utilized to detect antibodies which specifically bind to zsig15 proteins or peptides. Exemplary assays are described in detail in Antibodies: A Laboratory Manual, Harlow and Lane (Eds.), Cold Spring Harbor Laboratory Press, 1988. Representative examples

of such assays include: concurrent immunoelectrophoresis, radioimmunoassay, radioimmuno-precipitation, enzyme-linked immunosorbent assay (ELISA), dot blot or Western blot assay, inhibition or competition assay, and sandwich
5 assay. In addition, antibodies can be screened for binding to wild-type versus mutant zsig15 polypeptide.

Antibodies to zsig15 may be used for tagging cells that express zsig15 for isolating zsig15 by affinity purification; for diagnostic assays for
10 determining circulating levels of zsig15 polypeptides; for detecting or quantitating soluble zsig15 as marker of underlying pathology or disease; in analytical methods employing FACS; for screening expression libraries; for generating anti-idiotypic antibodies; and as neutralizing
15 antibodies or as antagonists to block zsig15 activity in vitro and in vivo. Suitable direct tags or labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent markers, chemiluminescent markers, magnetic particles and the like; indirect tags
20 or labels may feature use of biotin-avidin or other complement/anti-complement pairs as intermediates. Antibodies herein may also be directly or indirectly conjugated to drugs, toxins, radionuclides and the like, and these conjugates used for in vivo diagnostic or
25 therapeutic applications. Moreover, antibodies to zsig15 or fragments thereof may be used in vitro to detect denatured zsig15 or fragments thereof in assays, for example, Western Blots or other assays known in the art.

Antibodies or polypeptides herein can also be
30 directly or indirectly conjugated another moiety, for example, to drugs, toxins, radionuclides and the like, and these conjugates used for in vivo diagnostic or therapeutic applications. For instance, polypeptides or antibodies of the present invention can be used to

identify or treat tissues or organs that express a corresponding anti-complementary molecule (receptor or antigen, respectively, for instance). More specifically, zsig15 polypeptides or anti-zsig15 antibodies, or
5 bioactive fragments or portions thereof, can be coupled to detectable or cytotoxic molecules and delivered to a mammal having cells, tissues or organs that express the anti-complementary molecule.

Suitable detectable moieties may be directly or
10 indirectly attached to the polypeptide or antibody, and include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent markers, chemiluminescent markers, magnetic particles and the like. Suitable cytotoxic molecules may be directly or indirectly
15 attached to the polypeptide or antibody, and include bacterial or plant toxins (for instance, diphtheria toxin, *Pseudomonas* exotoxin, ricin, abrin and the like), as well as therapeutic radionuclides, such as iodine-131, rhenium-188 or yttrium-90 (either directly attached to
20 the polypeptide or antibody, or indirectly attached through means of a chelating moiety, for instance). Polypeptides or antibodies may also be conjugated to cytotoxic drugs, such as adriamycin. For indirect attachment of a detectable or cytotoxic molecule, the
25 detectable or cytotoxic molecule can be conjugated with a member of a complementary/ anticomplementary pair, where the other member is bound to the polypeptide or antibody portion. For these purposes, biotin/streptavidin is an exemplary complementary/ anticomplementary pair.

30 In another embodiment, polypeptide-toxin fusion proteins or antibody-toxin fusion proteins can be used for targeted cell or tissue inhibition or ablation (for instance, to treat cancer cells or tissues). Alternatively, if the polypeptide has multiple functional
35 domains (i.e., an activation domain or a ligand binding domain, plus a targeting domain), a fusion protein

including only the targeting domain may be suitable for directing a detectable molecule, a cytotoxic molecule or a complementary molecule to a cell or tissue type of interest. In instances where the domain only fusion protein includes a complementary molecule, the anti-complementary molecule can be conjugated to a detectable or cytotoxic molecule. Such domain-complementary molecule fusion proteins thus represent a generic targeting vehicle for cell/tissue-specific delivery of generic anti-complementary-detectable/ cytotoxic molecule conjugates.

In another embodiment, zsig15-cytokine fusion proteins or antibody-cytokine fusion proteins can be used for enhancing *in vivo* killing of target tissues (for example, blood, colon, breast and bone marrow cancers), if the zsig15 polypeptide or anti-zsig15 antibody targets the hyperproliferative cell (See, generally, Hornick et al., Blood 89:4437-47, 1997). They described fusion proteins enable targeting of a cytokine to a desired site of action, thereby providing an elevated local concentration of cytokine. Suitable zsig15 polypeptides or anti-zsig15 antibodies target an undesirable cell or tissue (i.e., a tumor or a leukemia), and the fused cytokine mediated improved target cell lysis by effector cells. Suitable cytokines for this purpose include interleukin 2 and granulocyte-macrophage colony-stimulating factor (GM-CSF), for instance.

The bioactive polypeptide or antibody conjugates described herein can be delivered intravenously, intraarterially or intraductally, or may be introduced locally at the intended site of action.

Molecules of the present invention can be used to identify and isolate receptors involved in cancer metastases. For example, proteins and peptides of the present invention can be immobilized on a column and membrane preparations run over the column (Immobilized

Affinity Ligand Techniques, Hermanson et al., eds., Academic Press, San Diego, CA, 1992, pp.195-202). Proteins and peptides can also be radiolabeled (Methods in Enzymol., vol. 182, "Guide to Protein Purification", 5 M. Deutscher, ed., Acad. Press, San Diego, 1990, 721-737) or photoaffinity labeled (Brunner et al., Ann. Rev. Biochem. 62:483-514, 1993 and Fedan et al., Biochem. Pharmacol. 33:1167-1180, 1984) and specific cell-surface proteins can be identified.

10 The polypeptides, nucleic acid and/or antibodies of the present invention can be used in treatment of disorders associated with prostate, colon or breast cancer. The molecules of the present invention can be used to modulate or to treat or prevent
15 development of pathological conditions. In particular, certain cancers, inflammatory and hyperplastic diseases may be amenable to such diagnosis, treatment or prevention.

 Polynucleotides encoding zsig15 polypeptides
20 are useful within gene therapy applications where it is desired to increase or inhibit zsig15 activity. If a mammal has a mutated or absent zsig15 gene, the zsig15 gene can be introduced into the cells of the mammal. In one embodiment, a gene encoding a zsig15 polypeptide is
25 introduced in vivo in a viral vector. Such vectors include an attenuated or defective DNA virus, such as, but not limited to, herpes simplex virus (HSV), papillomavirus, Epstein Barr virus (EBV), adenovirus, adeno-associated virus (AAV), and the like. Defective
30 viruses, which entirely or almost entirely lack viral genes, are preferred. A defective virus is not infective after introduction into a cell. Use of defective viral vectors allows for administration to cells in a specific, localized area, without concern that the vector can
35 infect other cells. Examples of particular vectors include, but are not limited to, a defective herpes

simplex virus 1 (HSV1) vector (Kaplitt et al., Molec. Cell. Neurosci. 2:320-30, 1991); an attenuated adenovirus vector, such as the vector described by Stratford-Perricaudet et al., J. Clin. Invest. 90:626-30, 1992; and
5 a defective adeno-associated virus vector (Samulski et al., J. Virol. 61:3096-101, 1987; Samulski et al., J. Virol. 63:3822-8, 1989).

In another embodiment, a zsig15 gene can be introduced in a retroviral vector, e.g., as described in
10 Anderson et al., U.S. Patent No. 5,399,346; Mann et al. Cell 33:153, 1983; Temin et al., U.S. Patent No. 4,650,764; Temin et al., U.S. Patent No. 4,980,289; Markowitz et al., J. Virol. 62:1120, 1988; Temin et al., U.S. Patent No. 5,124,263; International Patent
15 Publication No. WO 95/07358, published March 16, 1995 by Dougherty et al.; and Kuo et al., Blood 82:845, 1993. Alternatively, the vector can be introduced by lipofection in vivo using liposomes. Synthetic cationic lipids can be used to prepare liposomes for in vivo
20 transfection of a gene encoding a marker (Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7, 1987; Mackey et al., Proc. Natl. Acad. Sci. USA 85:8027-31, 1988). The use of lipofection to introduce exogenous genes into specific organs in vivo has certain practical advantages.
25 Molecular targeting of liposomes to specific cells represents one area of benefit. More particularly, directing transfection to particular cells represents one area of benefit. For instance, directing transfection to particular cell types would be particularly advantageous
30 in a tissue with cellular heterogeneity, such as the pancreas, liver, kidney, and brain. Lipids may be chemically coupled to other molecules for the purpose of targeting. Targeted peptides (e.g., hormones or neurotransmitters), proteins such as antibodies, or non-
35 peptide molecules can be coupled to liposomes chemically.

It is possible to remove the target cells from the body; to introduce the vector as a naked DNA plasmid; and then to re-implant the transformed cells into the body. Naked DNA vectors for gene therapy can be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, use of a gene gun or use of a DNA vector transporter. See, e.g., Wu et al., J. Biol. Chem. 267:963-7, 1992; Wu et al., J. Biol. Chem. 263:14621-4, 1988.

Antisense methodology can be used to inhibit zsig15 gene transcription, such as to inhibit cell proliferation in vivo. Polynucleotides that are complementary to a segment of a zsig15-encoding polynucleotide (e.g., a polynucleotide as set forth in SEQ ID NO:1) are designed to bind to zsig15-encoding mRNA and to inhibit translation of such mRNA. Such antisense polynucleotides are used to inhibit expression of zsig15 polypeptide-encoding genes in cell culture or in a subject.

The present invention also provides reagents which will find use in diagnostic applications. For example, the zsig15 gene, a probe comprising zsig15 DNA or RNA or a subsequence thereof can be used to determine if the zsig15 gene is present on chromosome 19 or if a mutation has occurred. Detectable chromosomal aberrations at the zsig15 gene locus include, but are not limited to, aneuploidy, gene copy number changes, insertions, deletions, restriction site changes and rearrangements. Such aberrations can be detected using polynucleotides of the present invention by employing molecular genetic techniques, such as restriction fragment length polymorphism (RFLP) analysis, short tandem repeat (STR) analysis employing PCR techniques, and other genetic linkage analysis techniques known in

the art (Sambrook et al., ibid.; Ausubel et. al., ibid.; Marian, Chest 108:255-65, 1995).

Transgenic mice, engineered to express the zsig15 gene, and mice that exhibit a complete absence of zsig15 gene function, referred to as "knockout mice" (Snouwaert et al., Science 257:1083, 1992), may also be generated (Lowell et al., Nature 366:740-42, 1993). These mice may be employed to study the zsig15 gene and the protein encoded thereby in an *in vivo* system.

Radiation hybrid mapping is a somatic cell genetic technique developed for constructing high-resolution, contiguous maps of mammalian chromosomes (Cox et al., Science 250:245-50, 1990). Partial or full knowledge of a gene's sequence allows one to design PCR primers suitable for use with chromosomal radiation hybrid mapping panels. Radiation hybrid mapping panels are commercially available which cover the entire human genome, such as the Stanford G3 RH Panel and the GeneBridge 4 RH Panel (Research Genetics, Inc., Huntsville, AL). These panels enable rapid, PCR-based chromosomal localizations and ordering of genes, sequence-tagged sites (STSs), and other nonpolymorphic and polymorphic markers within a region of interest. This includes establishing directly proportional physical distances between newly discovered genes of interest and previously mapped markers. The precise knowledge of a gene's position can be useful for a number of purposes, including: 1) determining if a sequence is part of an existing contig and obtaining additional surrounding genetic sequences in various forms, such as YACs, BACs or cDNA clones; 2) providing a possible candidate gene for an inheritable disease which shows linkage to the same chromosomal region; and 3) cross-referencing model organisms, such as mouse, which may aid in determining what function a particular gene might have.

Sequence tagged sites (STSs) can also be used independently for chromosomal localization. An STS is a DNA sequence that is unique in the human genome and can be used as a reference point for a particular chromosome or region of a chromosome. An STS is defined by a pair of oligonucleotide primers that are used in a polymerase chain reaction to specifically detect this site in the presence of all other genomic sequences. Since STSs are based solely on DNA sequence they can be completely described within an electronic database, for example, Database of Sequence Tagged Sites (dbSTS), GenBank, (National Center for Biological Information, National Institutes of Health, Bethesda, MD <http://www.ncbi.nlm.nih.gov>), and can be searched with a gene sequence of interest for the mapping data contained within these short genomic landmark STS sequences.

For pharmaceutical use, the proteins of the present invention are formulated for parenteral, particularly intravenous or subcutaneous, delivery according to conventional methods. Intravenous administration will be by bolus injection or infusion over a typical period of one to several hours. In general, pharmaceutical formulations will include a zsig15 protein in combination with a pharmaceutically acceptable vehicle, such as saline, buffered saline, 5% dextrose in water or the like. Formulations may further include one or more excipients, preservatives, solubilizers, buffering agents, albumin to prevent protein loss on vial surfaces, etc. Methods of formulation are well known in the art and are disclosed, for example, in Remington: The Science and Practice of Pharmacy, Gennaro, ed., Mack Publishing Co., Easton, PA, 19th ed., 1995. Therapeutic doses will generally be in the range of 0.1 to 100 µg/kg of patient weight per day, preferably 0.5-20 µg/kg per day, with the exact dose determined by the clinician according to accepted

standards, taking into account the nature and severity of the condition to be treated, patient traits, etc. Determination of dose is within the level of ordinary skill in the art. The proteins may be administered for
5 acute treatment, over one week or less, often over a period of one to three days or may be used in chronic treatment, over several months or years.

The invention is further illustrated by the following non-limiting examples.

10

EXAMPLES

Example 1

Scanning of a cDNA database for cDNAs
15 containing a secretion sequence revealed an expressed sequence tag (EST) that was novel and did not have homology to any known protein. The cDNA is from a human colon cDNA library.

Confirmation of the EST sequence was made by
20 excising the cDNA from its plasmid using a *Sal* I (nucleotide 1 of SEQ ID NO: 1) to Not I (nucleotide 1726 of SEQ ID NO: 1) and doing a sequence analyses of the cDNA from which the EST originated. The analyses revealed that the cDNA encompassed the entire coding
25 region of the DNA encoding zsig15.

Example 2

Northerns were performed using Human Multiple Tissue Blots and Human RNA Master dot blots from Clontech
30 (Palo Alto, CA). The probe was approximately 30 bp oligonucleotide ZC11366 (SEQ ID NO: 3). The probe was end labeled using T4 Polynucleotide Kinase (Life Technologies, Inc., Gaithersburg, MD) and T4 Polynucleotide Kinase Forward Buffer (Life Technologies,
35 Inc.). The probe was purified using a NUCTRAP push columns (Stratagene, La Jolla, CA). EXPRESSHYB

(Clontech) solution was used for prehybridization and as a hybridizing solution for the Northern blots. Hybridization took place at 42°C, and the blots were washed in 2X SSC and 0.1% SDS at RT, followed by a wash
 5 in 1 X SSC and 0.1% SDS at 50°C. After an overnight exposure a third wash at 65°C in 1X SSC. 0.1% SDS was done. Two transcripts were observed at 1.7 and 2.5 kb with a strong signal in prostate, with weaker signals seen in bone marrow and pancreas, colon mucosal lining
 10 and spinal cord. Dot blots revealed positive signals for prostate and bone marrow.

Tumor blots were probed with the insert from clone described previously. The probe was labeled as described previously and the tumor blots were washed at
 15 2X SSC, 0.1% SDS at RT, followed by a wash at 65°C in 0.1X SSC, 0.1% SDS. The human colon tumor blot (Invitrogen, Carlsbad, CA) contained total RNA isolated from human tumor and normal tissues from four different donors, where the normal/tumor pair was excised from the
 20 same operational site. Table 5 summarizes the results.

Table 5

sex	age	tumor description	normal expression level	tumor expression level
M	37	adeno- carcinoma	++++	-
M	59	adeno- carcinoma	+++	-
M	33	differentiated adeno- carcinoma	+++	++
M	56	poorly- moderately differentiated adenocarcinoma	-	++++

The human breast tumor blot (Invitrogen) contained total RNA isolated from human tumor and normal tissues from four different donors, where the normal/tumor pair was excised from the same operational
5 site. Table 6 summarizes the results.

Table 6

sex	age	tumor description	normal expression level	tumor expression level
F	57	invasive ductal carcinoma	++	-
F	50	invasive ductal carcinoma	+	-
M	51	invasive ductal carcinoma	-	+++
M	48	invasive ductal carcinoma	++	-

These data suggest that zsig15 mRNA is expressed at levels that vary with the cell's or tumor's
10 differentiated state.

Example 3

Zsig15 was mapped to chromosome 19 using the
15 commercially available "GeneBridge 4 Radiation Hybrid Panel" (Research Genetics, Inc., Huntsville, AL). The GeneBridge 4 Radiation Hybrid Panel contains DNAs from each of 93 radiation hybrid clones, plus two control DNAs (the HFL donor and the A23 recipient). A publicly
20 available WWW server (<http://www-genome.wi.mit.edu/cgi-bin/contig/rhmapper.pl>) allows mapping relative to the Whitehead Institute/MIT Center for Genome Research's

radiation hybrid map of the human genome (the "WICGR" radiation hybrid map) which was constructed with the GeneBridge 4 Radiation Hybrid Panel.

For the mapping of Zsig15 with the "GeneBridge 4 RH Panel", 25 μ l reactions were set up in a 96-well microtiter plate (Stratagene, La Jolla, CA) and used in a "RoboCycler Gradient 96" thermal cycler (Stratagene). Each of the 95 PCR reactions consisted of 2.5 μ l "10X KlenTaq reaction buffer" (Clontech Laboratories, Inc., Palo Alto, CA), 2 μ l dNTPs mix (2.5 mM each, Perkin-Elmer, Foster City, CA), 1.25 μ l sense primer, ZC 11,369, 5' CGG CAA TGG ACC CCT AAG AA 3', 1.25 μ l antisense primer, ZC 12,162, (SEQ ID NO:) 5' TCC TCC TGG CGG CAC ACG AA 3', 2.5 μ l "RediLoad" (Research Genetics, Inc., Huntsville, AL), 0.5 μ l "50X Advantage KlenTaq Polymerase Mix" (Clontech Laboratories, Inc.), 25 ng of DNA from an individual hybrid clone or control and ddH₂O for a total volume of 25 μ l. The reactions were overlaid with an equal amount of mineral oil and sealed. The PCR cycler conditions were as follows: an initial 1 cycle 5 minute denaturation at 95°C, 35 cycles of a 1 minute denaturation at 95°C, 1 minute annealing at 70°C and 1.5 minute extension at 72°C, followed by a final 1 cycle extension of 7 minutes at 72°C. The reactions were separated by electrophoresis on a 3% NuSieve GTG agarose gel (FMC Bioproducts, Rockland, ME).

The results showed that Zsig15 maps 323.29 cR_3000 from the top of the human chromosome 19 linkage group on the WICGR radiation hybrid map. Proximal and distal framework markers were D19S827 and D19S420, respectively. These markers position Zsig15 in the 19q13.2 region on the integrated LDB chromosome 19 map (The Genetic Location Database, University of Southampton, WWW server: [http:// cedar.genetics.soton.ac.uk/public_html/](http://cedar.genetics.soton.ac.uk/public_html/)).

Example 4

Creation of mammalian expression vectors zsig15NF/pZP9, zsig15CF/pZP9, soluble zsig15sNF/pZP9, soluble zsig15sCF/pZP9 and zsig15/pZP9

5 Five expression vectors were prepared for the zsig15 polypeptide, zsig15CF/pZP9 and zsig15NF/pZP9 were designed to express a full length zsig15 polypeptide with a C- or N-terminal FLAG tag (SEQ ID NO:5), zsig15sNF/pZP9, zsig15sCF/pZP9 were designed to express a
10 soluble zsig15 polypeptide having an N- or C-terminal FLAG tag, and zsig15/pZP9, was designed to express an untagged zsig15 polypeptide.

ZSIG15/pZP9

An approximately 1726 bp restriction digest
15 fragment of zsig-15 DNA was derived from the clone described in Example 1 above by restriction digest with enzymes Sal I and Not I. The resultant restriction fragment was visualized by agarose gel electrophoresis. A band of the predicted size was excised and the DNA was
20 purified from the gel with a QIAQUICK[®] column (Qiagen) according to the manufacturer's instructions.

The excised, restriction digested zsig15 DNA fragment was subcloned into plasmid pZP9 which had been cut with Xho I and Not I. Plasmid pZP9 (deposited at the
25 American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD) is a mammalian expression vector containing an expression cassette having the mouse metallothionein-1 promoter, multiple restriction sites for insertion of coding sequences, a stop codon and a
30 human growth hormone terminator. The plasmid also has an *E. coli* origin of replication, a mammalian selectable marker expression unit having an SV40 promoter, enhancer and origin of replication, a DHFR gene and the SV40 terminator.

35 The insert and expression vector were ligated overnight at 16°C, followed by electroporation into DH10B competent cells (GIBCO BRL, Gaithersburg, MD) according

to manufacturer's direction. The cells were plated onto LB plates containing 50 mg/ml ampicillin, and incubated at 37°C overnight. Colonies were screened by PCR using oligonucleotide primers to the vector, ZC13006 (SEQ ID NO:6) and ZC13007 (SEQ ID NO:7). The insert sequence of positive clones was verified by sequence analysis. A large scale plasmid preparation was done using a QIAGEN® Maxi prep kit (Qiagen) according to manufacturer's instructions.

10

zSIG25CF/pZP9

A 1.3 kb PCR generated zsig15 DNA fragment was created using two rounds of PCR to remove a Bam HI site (nucleotide 423 of SEQ ID NO:1 changed from C to A). Oligonucleotide primers ZC 13809 (SEQ ID NO:8) and ZC13812 (SEQ ID NO:9) were used to generate a ~392 bp 5' fragment. Oligonucleotide primers ZC 13804 (SEQ ID NO:10) and ZC13815 (SEQ ID NO:11) were used to generate a ~925 bp overlapping 3' fragment. Zsig15/pZP9 as described above, was used as a template, with an initial denaturation at 94°C for 1 minute 30 seconds, 12 cycles of 94°C, 10 seconds, 76°C 20 seconds, 72°C for 1 minute 30 seconds, followed by a 10 minute extension at 72°C. The resulting fragments were purified using a S-300 MicroSpin column (Pharmacia LKB Biotechnology Inc., Piscataway, NJ) according the manufacturer's instructions. A second round of PCR was done to join the 5' and 3' fragments using oligonucleotide primers ZC 13809 (SEQ ID NO:8) and ZC13815 (SEQ ID NO:11). The 5' and 3' first round fragments were used as templates for the second round of PCR, with an initial denaturation at 94°C for 1 minute 30 seconds, 12 cycles of 94°C, 10 seconds, 76°C 20 seconds, 72°C for 1 minute 30 seconds, followed by a 10 minute extension at 72°C. The resulting 1.3 kb second round fragment visualized by agarose gel

electrophoresis, excised and gel purified using QIAQUICK[®] column (Qiagen, Inc., Chatsworth, CA) according to manufacturer's instruction.

The purified PCR fragment was digested with the
5 restriction enzymes Bam HI and Eco RI, visualized by
agarose gel electrophoresis, excised and gel purified
using QIAQUICK[®] column (Qiagen) as above. The
restriction fragment was ligated into a Eco RI-Bam HI
restriction digested CF/pZP9 expression vector. The
10 zsig15/CFpZP9 expression vector uses the native zsig15
signal peptide, and the FLAG epitope (SEQ ID NO:5) is
attached at the C-terminus as a purification aid.
Plasmid CF/pZP9 (deposited at the American Type Culture
Collection, 12301 Parklawn Drive, Rockville, MD) is a
15 mammalian expression vector containing an expression
cassette having the mouse metallothionein-1 promoter,
multiple restriction sites for insertion of coding
sequences, a sequence encoding the FLAG tag (SEQ ID
NO:5), a stop codon and a human growth hormone
20 terminator. The plasmid also has an *E. coli* origin of
replication, a mammalian selectable marker expression
unit having an SV40 promoter, enhancer and origin of
replication, a DHFR gene and the SV40 terminator.

25 zSIG25NF/pZP9

A approximately 1.26 kb fragment of ZSIG-15 DNA
was derived by two rounds of PCR as described above,
removing a Bam HI site by changing nucleotide 423 of SEQ
ID NO:1 from C to A and adding a stop codon at the 3'
30 end. In the first round, oligonucleotides ZC13811 (SEQ
ID NO:12) and ZC13812 (SEQ ID NO:9) were used, with
zsig15/pZP9 as a template, generating an approximately
345 bp 5' fragment. Oligonucleotide primers ZC13804 (SEQ
ID NO:10) and ZC13810 (SEQ ID NO:13) were used to
35 generate an approximately 925 bp overlapping 3' fragment.
The 5' and 3' first round fragments were used as

templates for the second round PCR with oligonucleotide primers ZC13811 (SEQ ID NO:12) and ZC13810 (SEQ ID NO:13). The resulting 1.26 kb second round fragment was visualized by agarose gel electrophoresis, excised and
5 gel purified using QIAQUICK[®] column (Qiagen, Inc., Chatsworth, CA) according to manufacturer's instruction.

The purified PCR fragment was digested with the restriction enzymes Bam HI and Xba I and ligated into a Bam HI/Xba I restriction digested NFpZP9 expression
10 vector as described above. The zsig15NF/pZP9 vector incorporates the TPA leader and attaches the FLAG tag (SEQ ID NO:5) to the N-terminal of the zsig15 polypeptide-encoding polynucleotide sequence. Plasmid NF/pZP9 (deposited at the American Type Culture
15 Collection, 12301 Parklawn Drive, Rockville, MD) is a mammalian expression vector containing an expression cassette having the mouse metallothionein-1 promoter, a TPA leader peptide followed by the sequence encoding the FLAG tag (SEQ ID NO:5), multiple restriction sites for
20 insertion of coding sequences, and a human growth hormone terminator. The plasmid also contains an *E. coli* origin of replication, a mammalian selectable marker expression unit having an SV40 promoter, enhancer and origin of replication, a DHFR gene and the SV40 terminator.

25

zsig15sCF/pZP9

A approximately 1.18 kb soluble ZSIG-15 DNA fragment was derived by two rounds of PCR as described above, removing a Bam HI site by changing nucleotide 423
30 of SEQ ID NO:1 from C to A and truncating the sequence at amino acid residue 394 (Glu) of SEQ ID NO:2 just prior to the hydrophobic region. In the first round, oligonucleotides ZC13809 (SEQ ID NO:8) and ZC13812 (SEQ ID NO:9) were used, with zsig15/pZP9 as a template,
35 generating an approximately 392 bp 5' fragment. Oligonucleotide primers ZC13804 (SEQ ID NO:10) and

ZC13814 (SEQ ID NO:14) were used to generate an approximately 780 bp overlapping 3' fragment. The 5' and 3' first round fragments were used as templates for the second round PCR with oligonucleotide primers ZC13809 (SEQ ID NO:8) and ZC13814 (SEQ ID NO:14). The resulting 1.18 kb second round fragment was visualized by agarose gel electrophoresis, excised and gel purified using QIAQUICK[®] column (Qiagen, Inc., Chatsworth, CA) according to manufacturer's instruction.

10

zsig15sNF/pZP9

A approximately 1.14 kb soluble ZSIG-15 DNA fragment was derived by two rounds of PCR as described above, removing a Bam HI site by changing nucleotide 423 of SEQ ID NO:1 from C to A and truncating the sequence at amino acid residue 394 (Glu) of SEQ ID NO:2 just prior to the hydrophobic region and adding a stop codon at the 3' end. In the first round, oligonucleotides ZC13811 (SEQ ID NO:12) and ZC13812 (SEQ ID NO:9) were used, with zsig15/pZP9 as a template, to generate an approximately 345 bp 5' fragment. Oligonucleotide primers ZC13804 (SEQ ID NO:10) and ZC13813 (SEQ ID NO:15) were used to generate an approximately 780 bp overlapping 3' fragment. The 5' and 3' first round fragments were used as templates for the second round PCR with oligonucleotide primers ZC13813 (SEQ ID NO:15) and ZC13811 (SEQ ID NO:12). The resulting approximately 1.14 kb second round fragment was visualized by agarose gel electrophoresis, excised and gel purified using QIAQUICK[®] column (Qiagen, Inc.) according to manufacturer's instruction.

Approximately 20-30 ng of each of the restriction digested inserts and 0.05-0.06 pm of the corresponding vectors were independently ligated overnight at 16°C. 1.2 µl of each ligation reaction was independently electroporated into 38 µl of DH10B competent cells (GIBCO BRL, Gaithersburg, MD) according

to manufacturer's direction and plated onto LB plates containing 50 mg/ml ampicillin, and incubated overnight. Colonies were screened by PCR using oligonucleotide primers to the expression vectors, ZC13006 (SEQ ID NO:6) and ZC13007 (SEQ ID NO:7). The insert sequence of positive clones were verified by sequence analysis. A large scale plasmid preparation was done using a QIAGEN[®] Maxi prep kit (Qiagen) according to manufacturer's instructions.

10

Example 5

Mammalian Expression of zsig15

BHK 570 cells (ATCC NO: CRL-10314) were plated in 10 cm tissue culture dishes and allowed to grow to approximately 50 to 70% confluency overnight at 37°C, 5% CO₂, in DMEM/FBS media (DMEM, Gibco/BRL High Glucose, (Gibco BRL, Gaithersburg, MD), 5% fetal bovine serum (Hyclone, Logan, UT), 1 µM L-glutamine (JRH Biosciences, Lenexa, KS), 1 µM sodium pyruvate (Gibco BRL)). The cells were then transfected with the plasmid zsig15NF/pZP9 (full length N-terminal FLAG tag), zsig15CF/pZP9 (full length C-terminal FLAG tag), zsig15sNF/pZP9 (soluble N-terminal FLAG tag) and zsig15sCF/pZP9 (soluble C-terminal FLAG tag), using Lipofectamine[™] (Gibco BRL), in serum free (SF) media formulation (DMEM, 10 mg/ml transferrin, 5 mg/ml insulin, 2 mg/ml fetuin, 1% L-glutamine and 1% sodium pyruvate). Sixteen micrograms of each expression construct were separately diluted into 15 ml tubes to a total final volume of 640 µl with SF media. In separate tubes, 35 µl of Lipofectamine[™] (Gibco BRL) was mixed with 605 µl of SF medium. The Lipofectamine[™] mix was added to the expression construct mix and allowed to incubate approximately 30 minutes at room temperature. Five milliliters of SF media was added to the

DNA:LipofectamineTM mixture. Three plates of cells were rinsed once with 5 ml of SF media, aspirated, and the DNA:LipofectamineTM mixture was added. The cells were incubated at 37°C for five hours, then 6.4 ml of DMEM/10%
5 FBS, 1% PSN media was added to each plate. The plates were incubated at 37°C overnight and the DNA:LipofectamineTM mixture was replaced with fresh FBS/DMEM media the next day. On day 2 post-transfection, the cells were split into the selection media (DMEM/FBS
10 media from above with the addition of 1 µM methotrexate (Sigma Chemical Co., St. Louis, Mo.)) in 150 mm plates at 1:10, 1:20 and 1:50. The cells were refed at day 5 post-transfection with fresh selection media.

Approximately 10-12 days post-transfection, two
15 150 mm culture dishes of methotrexate resistant colonies were chosen, the media aspirated, the plates washed with 10 ml serum-free ESTEP 2 media (668.7g/50L DMEM (Gibco), 5.5 g/50L pyruvic acid, sodium salt 96% (Mallinckrodt), 185.0 g/50L NaHCO₃ (Mallinckrodt), 5.0 mg/ml, 25 ml/50L
20 insulin, 10.0 mg/ml and 25 ml/50 L transferrin). The wash media was aspirated and replaced with 5 ml serum-free ESTEP 2. Sterile Teflon mesh (Spectrum Medical Industries, Los Angeles, CA) pre-soaked in serum-free ESTEP 2 was then placed over the cells. A sterile
25 nitrocellulose filter pre-soaked in serum-free ESTEP 2 was then placed over the mesh. Orientation marks on the nitrocellulose were transferred to the culture dish. The plates were then incubated for 5-6 hours in a 37°C, 5% CO₂ incubator. Following incubation, the filter was
30 removed, and the media aspirated and replaced with DMEM/5% FBS, 1X PSN (Gibco BRL) media. The filters were blocked in 2.5% nonfat dry milk/Western A buffer (Western A: 50mM Tris pH 7.4, 5 mM EDTA, 0.05% NP-40, 150 mM NaCl and 0.25% gelatin) overnight at 4°C on a rotating shaker.
35 The filter was then incubated with a goat anti-human

FLAG-HRP conjugate at a 1:4000 dilution (5 μ l antibody in 20 ml buffer) in 2.5% nonfat dry milk/Western A buffer (Western A: 50mM Tris pH 7.4, 5 mM EDTA, 0.05% NP-40, 150 mM NaCl and 0.25% gelatin) at room temperature for 1 hour
5 on a rotating shaker. The filter was then washed three times at room temperature in PBS plus 0.1% Tween 20, 15 minutes per wash. The filter was developed with ECL reagent (Amersham Corp., Arlington Heights, IL) according to the manufacturer's directions and exposed to film
10 (Hyperfilm ECL, Amersham) for approximately 5 minutes.

The film was aligned with the plate containing the colonies. Using the film as a guide, suitable colonies were selected. Sterile, 3 mm colonizing discs (PGC Scientific Corp., Frederick, MD) were soaked in
15 trypsin, and placed on the colonies. The colonies were transferred into 200 μ l of selection medium in a 96 well plate. A series of seven, two-fold dilutions were carried out for each colony. The 150 mm culture dish was then trypsinized and the remainder of the cells are
20 pooled and split into two T162 flasks containing DMEM/5% FBS and 1 μ M MTX media. The cells were grown for one week at 37°C at which time the wells which received the lowest dilution of cells which are now at the optimum density were selected, trypsinized and transferred to a
25 12 well plate containing selection media.

The clones were expanded directly from the 12 well plate to 2 T-75 flasks. One flask from each clone is grown in serum-free ESTEP 2 and the media harvested for Western Blot analysis. Clones of each of the expression
30 constructs, based on Western blot analysis were selected, pooled together and transferred to large scale culture.

Example 6

35 Large Scale Mammalian Expression of zsig15

One T-162 flask, containing confluent cells expressing zsig15s/NF and one flask containing zsig15s/CF expressing cells, obtained from the expression procedure described above, were expanded into five T-162 flasks.

- 5 One of the five resulting flasks was used to freeze down four cryovials, and the other four flasks were used to generate a Nunc cell factory.

The cells from the four T-162 flasks of zsig15s/NF and zsig15s/CF were combined and used to seed
10 two Nunc cell factories (10 layers, commercially available from VWR). Briefly, the cells from the T-162 flasks described above were detached using trypsin, pooled, and added to 1.5 liters ESTEP1 media (668.7g/50L DMEM (Gibco), 5.5 g/50L pyruvic acid, sodium salt 96%
15 (Mallinckrodt), 185.0 g/50L NaHCO₃ (Mallinckrodt), 5.0 mg/ml and 25 ml/50L insulin (JRH Biosciences), 10.0 mg/ml and 25 ml/50L transferrin (JRH Biosciences), 2.5L/50L fetal bovine serum (characterized) (Hyclone), 1 μ M MTX, with pH adjusted to 7.05 +/-0.05) prewarmed to 37°C. The
20 media containing the cells was then poured into the Nunc cell factories via a funnel. The cell factories were placed in a 37°C/5.0% CO₂ incubator.

At 80-100% confluence, a visual contamination test (phenol red color change) was performed on the Nunc
25 cell factories. Since no contamination was observed, supernatant from the confluent factories was poured into a small harvest container, sampled and discarded. The adherent cells were then washed once with 400 ml PBS. To detach the cells from the factories, 100 mls of trypsin
30 was added to each and removed and the cells were then incubated for 5 to 10 minutes in the residual trypsin. The cells were collected following two, 200 ml washes of ESTEP1 media. To each of ten ESTEP1 media-containing bottles (1.5 liters each, at 37°C) was added 40 mls of
35 collected cells. One 1.5 liter bottle was then used to

fill one Nunc factory. Each cell factory was placed in a 37°C/5.0% CO₂ incubator.

At 80-90% confluence, a visual contamination test (phenol red color change) was performed on the Nunc cell factories. Since no contamination was observed, supernatant from the confluent factories was poured into a small harvest container, sampled and discarded. Cells were then washed once with 400 ml PBS. 1.5 liters of ESTEP2 media (668.7g/50L DMEM (Gibco), 5.5 g/50L pyruvic acid, sodium salt 96% (Mallinckrodt), 185.0 g/50L NaHCO₃ (Mallinckrodt), 5.0 mg/ml, 25 ml/50L insulin, 10.0 mg/ml and 25 ml/50L transferrin) was added to each Nunc cell factory. The cell factories were incubated at 37°C/5.0% CO₂.

At approximately 40 hours a visual contamination test (phenol red color change) was performed on the Nunc cell factories. Supernatant from each factory was poured into small harvest containers. A total of 15 liters was collected from all 10 factories. Fresh serum-free media (1.5 liters) was poured into each Nunc cell factory, and the factories were incubated at 37°C/5.0% CO₂. One ml of supernatant harvest was transferred to a microscope slide, and subjected to microscopic analysis for contamination. The contents of the small harvest containers for each factory were pooled and immediately filtered. A second harvest was then performed, substantially as described above at 50 hours (15 L were obtained) and the cell factories were discarded thereafter. An aseptically assembled filter train apparatus was used for aseptic filtration of the harvest supernatant (conditioned media). Assembly was as follows: tubing was wire-tied to an Opti-Cap filter (Millipore Corp., Bedford, MA) and a Gelman Supercap 50 filter (Gelman Sciences, Ann Arbor, MI). The Supercap 50 filter was also attached to a sterile capped container located in a hood; tubing located upstream of the

Millipore Opti-cap filter was inserted into a peristaltic pump; and the free end of the tubing was placed in the large harvest container. The peristaltic pump was run between 200 and 300 rpm, until all of the conditioned media passed through the 0.22 μ m final filter into a sterile collection container. The filtrate was placed in a 4°C cold room pending purification. The media was concentrated 10X with a Millipore 5 kDA cut off concentrator (Millipore Corp., Bedford, MA) according to manufacturer's direction and subjected to Western Blot analysis using an anti-FLAG tag antibody (Kodak).

Example 7

Unless otherwise noted, all operations are carried out at 4°C. The following procedure is used to purify zsig15 containing N-terminal or C-terminal flag tags. A total of 25 liters of conditioned media from baby hamster kidney (BHK) cells is sequentially sterile filtered through a 4 inch, 0.2 μ m Millipore (Bedford, MA) OptiCap capsule filter and a 0.2 μ m Gelman (Ann Arbor, MI) Supercap 50. The material is concentrated to about 1.3 liters using an Amicon (Beverly, MA) DC 10L concentrator fitted with a 3000 kDa cutoff membrane. The concentrated material is sterile-filtered again with is be added to the concentrated conditioned media to final concentrations of 2.5 mM ethylenediaminetetraacetic acid (EDTA, Sigma Chemical Co. St. Louis, MO), 0.001 mM leupeptin (Boehringer-Mannheim, Indianapolis, IN), 0.001 mM pepstatin (Boehringer-Mannheim) and 0.4 mM Pefabloc (Boehringer-Mannheim). A 25.0 ml sample of anti-Flag Sepharose (Eastman Kodak, Rochester, NY) is added to the sample for batch adsorption and the mixture is gently agitated on a Wheaton (Millville, NJ) roller culture apparatus for 18.0 h at 4°C.

The mixture is then poured into a 5.0 x 20.0 cm Econo-Column (Bio-Rad, Laboratories, Hercules, CA) and

the gel is washed with 30 column volumes of phosphate buffered saline (PBS). The unretained flow-through fraction is discarded. Once the absorbance of the effluent at 280 nM is less than 0.05, flow-through the column is reduced to zero and the anti-Flag Sepharose gel is washed batchwise with 2.0 column volumes of PBS containing 0.2 mg/ml of Flag peptide, N-AspTyrLysAspAspAspAspLys-C (Seq. ID. No. 5; Eastman Kodak). After 1.0 h at 4°C, flow is resumed and the eluted protein is collected. This fraction is referred to as the peptide elution. The anti-Flag Sepharose gel is washed with 2.0 column volumes of 0.1M glycine, pH 2.5, and the glycine wash is collected separately. The pH of the glycine-eluted fraction is adjusted to 7.0 by the addition of a small volume of 10X PBS and is stored at 4°C for future analysis if needed.

The peptide elution is concentrated to 5.0 ml using a 5,000 molecular weight cutoff membrane concentrator (Millipore, Bedford, MA), according to the manufacturer's instructions. The concentrated peptide elution is separated from free peptide by chromatography on a 1.5 x 50 cm Sephadex G-50 (Pharmacia, Piscataway, NJ) column equilibrated in PBS at a flow rate of 1.0 ml/min using a BioCad Sprint HPLC system (PerSeptive BioSystems, Framingham, MA). Two-ml fractions are collected and the absorbance at 280 nM is monitored. The first peak of material absorbing at 280 nM and eluting near the void volume of the column is collected. The purity of the zsig15 NF and zsig15 CF is monitored by SDS-PAGE and Western analysis with anti-Flag M2 antibodies (Kodak).

The protein concentration of the purified proteins is performed by BCA analysis (Pierce, Rockford, IL) and the material is aliquoted, and stored at -80°C.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have

been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended
5 claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: ZymoGenetics, Inc.
1201 Eastlake Avenue East
Seattle
WA
USA
98102
- (ii) TITLE OF THE INVENTION: NOVEL TUMOR ANTIGENS
- (iii) NUMBER OF SEQUENCES: 15
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: ZymoGenetics, Inc.
 - (B) STREET: 1201 Eastlake Avenue East
 - (C) CITY: Seattle
 - (D) STATE: WA
 - (E) COUNTRY: USA
 - (F) ZIP: 98102
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Sawislak, Deborah A
 - (B) REGISTRATION NUMBER: 37,438
 - (C) REFERENCE/DOCKET NUMBER: 97-14PC

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 206-442-6672
- (B) TELEFAX: 206-442-6678
- (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1733 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: Coding Sequence
- (B) LOCATION: 34...1344
- (D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GTCGACCCAC GCGTCCGACG CCACAGACGG GTC ATG AGC GCG GTA TTA CTG CTG	54
Met Ser Ala Val Leu Leu Leu	
1 5	
GCC CTC CTG GGG TTC ATC CTC CCA CTG CCA GGA GTG CAG GCG CTG CTC	102
Ala Leu Leu Gly Phe Ile Leu Pro Leu Pro Gly Val Gln Ala Leu Leu	
10 15 20	
TGC CAG TTT GGG ACA GTT CAG CTT GTG TGG AAG GTG TCC GAC CTA CCC	150
Cys Gln Phe Gly Thr Val Gln Leu Val Trp Lys Val Ser Asp Leu Pro	
25 30 35	
CGG CAA TGG ACC CCT AAG AAC ACC AGC TGC GAC AGC GGC TTG GGG TGC	198
Arg Gln Trp Thr Pro Lys Asn Thr Ser Cys Asp Ser Gly Leu Gly Cys	
40 45 50 55	
CAG GAC ACG TTG ATG CTC ATT GAG AGC GGA CCC CAA GTG AGC CTG GTG	246
Gln Asp Thr Leu Met Leu Ile Glu Ser Gly Pro Gln Val Ser Leu Val	
60 65 70	

CTC	TCC	AAG	GGC	TGC	ACG	GAG	GCC	AAG	GAC	CAG	GAG	CCC	CGC	GTC	ACT	294
Leu	Ser	Lys	Gly	Cys	Thr	Glu	Ala	Lys	Asp	Gln	Glu	Pro	Arg	Val	Thr	
			75					80					85			
GAG	CAC	CGG	ATG	GGC	CCC	GGC	CTC	TCC	CTG	ATC	TCC	TAC	ACC	TTC	GTG	342
Glu	His	Arg	Met	Gly	Pro	Gly	Leu	Ser	Leu	Ile	Ser	Tyr	Thr	Phe	Val	
		90					95					100				
TGC	CGC	CAG	GAG	GAC	TTC	TGC	AAC	AAC	CTC	GTT	AAC	TCC	CTC	CCG	CTT	390
Cys	Arg	Gln	Glu	Asp	Phe	Cys	Asn	Asn	Leu	Val	Asn	Ser	Leu	Pro	Leu	
	105					110					115					
TGG	GCC	CCA	CAG	CCC	CCA	GCA	GAC	CCA	GGA	TCC	TTG	AGG	TGC	CCA	GTC	438
Trp	Ala	Pro	Gln	Pro	Pro	Ala	Asp	Pro	Gly	Ser	Leu	Arg	Cys	Pro	Val	
120					125					130					135	
TGC	TTG	TCT	ATG	GAA	GGC	TGT	CTG	GAG	GGG	ACA	ACA	GAA	GAG	ATC	TGC	486
Cys	Leu	Ser	Met	Glu	Gly	Cys	Leu	Glu	Gly	Thr	Thr	Glu	Glu	Ile	Cys	
			140						145					150		
CCC	AAG	GGG	ACC	ACA	CAC	TGT	TAT	GAT	GGC	CTC	CTC	AGG	CTC	AGG	GGA	534
Pro	Lys	Gly	Thr	Thr	His	Cys	Tyr	Asp	Gly	Leu	Leu	Arg	Leu	Arg	Gly	
			155					160					165			
GGA	GGC	ATC	TTC	TCC	AAT	CTG	AGA	GTC	CAG	GGA	TGC	ATG	CCC	CAG	CCA	582
Gly	Gly	Ile	Phe	Ser	Asn	Leu	Arg	Val	Gln	Gly	Cys	Met	Pro	Gln	Pro	
		170					175					180				
GGT	TGC	AAC	CTG	CTC	AAT	GGG	ACA	CAG	GAA	ATT	GGG	CCC	GTG	GGT	ATG	630
Gly	Cys	Asn	Leu	Leu	Asn	Gly	Thr	Gln	Glu	Ile	Gly	Pro	Val	Gly	Met	
	185					190					195					
ACT	GAG	AAC	TGC	AAT	AGG	AAA	GAT	TTT	CTG	ACC	TGT	CAT	CGG	GGG	ACC	678
Thr	Glu	Asn	Cys	Asn	Arg	Lys	Asp	Phe	Leu	Thr	Cys	His	Arg	Gly	Thr	
200					205					210					215	
ACC	ATT	ATG	ACA	CAC	GGA	AAC	TTG	GCT	CAA	GAA	CCC	ACT	GAT	TGG	ACC	726
Thr	Ile	Met	Thr	His	Gly	Asn	Leu	Ala	Gln	Glu	Pro	Thr	Asp	Trp	Thr	
				220					225					230		
ACA	TCG	AAT	ACC	GAG	ATG	TGC	GAG	GTG	GGG	CAG	GTG	TGT	CAG	GAG	ACG	774
Thr	Ser	Asn	Thr	Glu	Met	Cys	Glu	Val	Gly	Gln	Val	Cys	Gln	Glu	Thr	
			235					240					245			

CTG CTG CTC ATA GAT GTA GGA CTC ACA TCA ACC CTG GTG GGG ACA AAA Leu Leu Leu Ile Asp Val Gly Leu Thr Ser Thr Leu Val Gly Thr Lys 250 255 260	822
GGC TGC AGC ACT GTT GGG GCT CAA AAT TCC CAG AAG ACC ACC ATC CAC Gly Cys Ser Thr Val Gly Ala Gln Asn Ser Gln Lys Thr Thr Ile His 265 270 275	870
TCA GCC CCT CCT GGG GTG CTT GTG GCC TCC TAT ACC CAC TTC TGC TCC Ser Ala Pro Pro Gly Val Leu Val Ala Ser Tyr Thr His Phe Cys Ser 280 285 290 295	918
TCG GAC CTG TGC AAT AGT GCC AGC AGC AGC AGC GTT CTG CTG AAC TCC Ser Asp Leu Cys Asn Ser Ala Ser Ser Ser Ser Val Leu Leu Asn Ser 300 305 310	966
CTC CCT CCT CAA GCT GCC CCT GTC CCA GGA GAC CGG CAG TGT CCT ACC Leu Pro Pro Gln Ala Ala Pro Val Pro Gly Asp Arg Gln Cys Pro Thr 315 320 325	1014
TGT GTG CAG CCC CTT GGA ACC TGT TCA AGT GGC TCC CCC CGA ATG ACC Cys Val Gln Pro Leu Gly Thr Cys Ser Ser Gly Ser Pro Arg Met Thr 330 335 340	1062
TGC CCC AGG GGC GCC ACT CAT TGT TAT GAT GGG TAC ATT CAT CTC TCA Cys Pro Arg Gly Ala Thr His Cys Tyr Asp Gly Tyr Ile His Leu Ser 345 350 355	1110
GGA GGT GGG CTG TCC ACC AAA ATG AGC ATT CAG GGC TGC GTG GCC CAA Gly Gly Gly Leu Ser Thr Lys Met Ser Ile Gln Gly Cys Val Ala Gln 360 365 370 375	1158
CCT TCC AGC TTC TTG TTG AAC CAC ACC AGA CAA ATC GGG ATC TTC TCT Pro Ser Ser Phe Leu Leu Asn His Thr Arg Gln Ile Gly Ile Phe Ser 380 385 390	1206
GCG CGT GAG AAG CGT GAT GTG CAG CCT CCT GCC TCT CAG CAT GAG GGA Ala Arg Glu Lys Arg Asp Val Gln Pro Pro Ala Ser Gln His Glu Gly 395 400 405	1254
GGT GGG GCT GAG GGC CTG GAG TCT CTC ACT TGG GGG GTG GGG CTG GCA Gly Gly Ala Glu Gly Leu Glu Ser Leu Thr Trp Gly Val Gly Leu Ala 410 415 420	1302

CTG GCC CCA GCG CTG TGG TGG AGA GTG GTT TGC CCT TCC TGC TAACTCTAT 1353
 Leu Ala Pro Ala Leu Trp Trp Arg Val Val Cys Pro Ser Cys
 425 430 435

TACCCCCACG ATTCTTCACC GCTGCTGACC ACCCACACTC AACCTCCCTC TGACCTCATA 1413
 ACCTAATGGC CTTGGACACC AGATTCTTTC CCATTCTGTC CATGAATCAT CTTCCCCACA 1473
 CACAATCATT CATATCTATT CACCTAACAG CAACACTGGG GAGAGCCTGG AGCATCCGGA 1533
 CTTGCCCTAT GGGAGAGGGG ACGCTGGAGG AGTGGCTGCA TGTATCTGAT AATACAGACC 1593
 CTGTCCTTTC TCCCAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 1653
 AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 1713
 AAAAAAAAAA GGGCGGCCGC 1733

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 437 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ser Ala Val Leu Leu Leu Ala Leu Leu Gly Phe Ile Leu Pro Leu
 1 5 10 15
 Pro Gly Val Gln Ala Leu Leu Cys Gln Phe Gly Thr Val Gln Leu Val
 20 25 30
 Trp Lys Val Ser Asp Leu Pro Arg Gln Trp Thr Pro Lys Asn Thr Ser
 35 40 45
 Cys Asp Ser Gly Leu Gly Cys Gln Asp Thr Leu Met Leu Ile Glu Ser
 50 55 60
 Gly Pro Gln Val Ser Leu Val Leu Ser Lys Gly Cys Thr Glu Ala Lys
 65 70 75 80
 Asp Gln Glu Pro Arg Val Thr Glu His Arg Met Gly Pro Gly Leu Ser
 85 90 95
 Leu Ile Ser Tyr Thr Phe Val Cys Arg Gln Glu Asp Phe Cys Asn Asn
 100 105 110
 Leu Val Asn Ser Leu Pro Leu Trp Ala Pro Gln Pro Pro Ala Asp Pro
 115 120 125
 Gly Ser Leu Arg Cys Pro Val Cys Leu Ser Met Glu Gly Cys Leu Glu
 130 135 140
 Gly Thr Thr Glu Glu Ile Cys Pro Lys Gly Thr Thr His Cys Tyr Asp
 145 150 155 160

Gly Leu Leu Arg Leu Arg Gly Gly Gly Ile Phe Ser Asn Leu Arg Val
 165 170 175
 Gln Gly Cys Met Pro Gln Pro Gly Cys Asn Leu Leu Asn Gly Thr Gln
 180 185 190
 Glu Ile Gly Pro Val Gly Met Thr Glu Asn Cys Asn Arg Lys Asp Phe
 195 200 205
 Leu Thr Cys His Arg Gly Thr Thr Ile Met Thr His Gly Asn Leu Ala
 210 215 220
 Gln Glu Pro Thr Asp Trp Thr Thr Ser Asn Thr Glu Met Cys Glu Val
 225 230 235 240
 Gly Gln Val Cys Gln Glu Thr Leu Leu Leu Ile Asp Val Gly Leu Thr
 245 250 255
 Ser Thr Leu Val Gly Thr Lys Gly Cys Ser Thr Val Gly Ala Gln Asn
 260 265 270
 Ser Gln Lys Thr Thr Ile His Ser Ala Pro Pro Gly Val Leu Val Ala
 275 280 285
 Ser Tyr Thr His Phe Cys Ser Ser Asp Leu Cys Asn Ser Ala Ser Ser
 290 295 300
 Ser Ser Val Leu Leu Asn Ser Leu Pro Pro Gln Ala Ala Pro Val Pro
 305 310 315 320
 Gly Asp Arg Gln Cys Pro Thr Cys Val Gln Pro Leu Gly Thr Cys Ser
 325 330 335
 Ser Gly Ser Pro Arg Met Thr Cys Pro Arg Gly Ala Thr His Cys Tyr
 340 345 350
 Asp Gly Tyr Ile His Leu Ser Gly Gly Gly Leu Ser Thr Lys Met Ser
 355 360 365
 Ile Gln Gly Cys Val Ala Gln Pro Ser Ser Phe Leu Leu Asn His Thr
 370 375 380
 Arg Gln Ile Gly Ile Phe Ser Ala Arg Glu Lys Arg Asp Val Gln Pro
 385 390 395 400
 Pro Ala Ser Gln His Glu Gly Gly Gly Ala Glu Gly Leu Glu Ser Leu
 405 410 415
 Thr Trp Gly Val Gly Leu Ala Leu Ala Pro Ala Leu Trp Trp Arg Val
 420 425 430
 Val Cys Pro Ser Cys
 435

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GCTGGTGTTC TTAGGGGTCC ATTGCCGGGG

30

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1311 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ATGWSNGCNG	TNYTNYTNYT	NGCNYTNYTN	GGNTTYATHY	TNCCNYTNCC	NGGNGTNCAR	60
GCNYTNYTNT	GYCARTTYGG	NACNGTNCAR	YTNGTNTGGA	ARGTNWSNGA	YYTNCCNMGN	120
CARTGGACNC	CNAARAAYAC	NWSNTGYGAY	WSNGGNYTNG	GNTGYCARGA	YACNYTNATG	180
YTNATHGARW	SNGGNCCNCA	RGTNWSNYTN	GTNYTNWSNA	ARGGNTGYAC	NGARGCNAAR	240
GAYCARGARC	CNMGNGTNAC	NGARCAYMGN	ATGGGNCCNG	GNYTNSNYT	NATHWSNTAY	300
ACNTTYGTNT	GYMGNCARGA	RGAYTTYTGY	AAYAAYYTNG	TNAAYSNYT	NCCNYTNTGG	360
GCNCCNCARC	CNCCNGCNGA	YCCNGGNWSN	YTNMGNTGYC	CNGTNTGYT	NWSNATGGAR	420
GGNTGYTNG	ARGGNACNAC	NGARGARATH	TGYCCNAARG	GNACNACNCA	YTGYTAYGAY	480
GGNYTNYTNM	GNYTNGMNGG	NGGNGGNATH	TTYWSNAAAY	TNMGNGTNCA	RGGNTGYATG	540
CCNCARCCNG	GNTGYAAYYT	NYTNAAAYGGN	ACNCARGARA	THGGNCCNGT	NGGNATGACN	600
GARAAYTGYA	AYMGNAARGA	YTTYTNAACN	TGYCAYMGNG	GNACNACNAT	HATGACNCAY	660
GGNAAYYTNG	CNCARGARCC	NACNGAYTGG	ACNACNWSNA	AYACNGARAT	GTGYGARGTN	720
GGNCARGTNT	GYCARGARAC	NYTNYTNYTN	ATHGAYGTNG	GNYTNAACNWS	NACNYTNGTN	780
GGNACNAARG	GNTGYWSNAC	NGTNGGNGCN	CARAAYWSNC	ARAARACNAC	NATHCAYWSN	840
GCNCCNCCNG	GNGTNYTNGT	NGCNWSNTAY	ACNCAYTTYT	GYWSNWSNGA	YYTNTGYAAY	900
WSNGCNWSNW	SNWSNWSNGT	NYTNYTNAAAY	WSNYTNCCNC	CNCARGCNGC	NCCNGTNCN	960
GGNGAYMGNC	ARTGYCCNAC	NTGYGTNCAR	CCNYTNGGNA	CNTGYWSNWS	NGGNWSNCCN	1020
MGNATGACNT	GYCCNMGNGG	NGCNACNCAY	TGYTAYGAYG	GNTAYATHCA	YYTNWSNGGN	1080
GGNGGNYTNW	SNACNAARAT	GWSNATHCAR	GGNTGYGTNG	CNCARCCNWS	NWSNTTYTNT	1140
YTNAAYCAYA	CNMGNCARAT	HGGNATHTTY	WSNGCNMGNG	ARAARMGNGA	YGTNCARCCN	1200
CCNGCNWSNC	ARCAYGARGG	NGGNGGNGCN	GARGGNYTNG	ARWSNYTNAC	NTGGGGNGTN	1260
GGNYTNGCNY	TNGCNCCNGC	NYTNTGGTGG	MNGTNGTNT	GYCCNWSNTG	Y	1311

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Asp Tyr Lys Asp Asp Asp Asp Lys
1 5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13006

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GGCTGTCCTC TAAGCGTCAC

20

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13007

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AGGGGTCACA GGGATGCCA

19

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13809

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TCGTAGAGAA TTCATGAGTG CGGTATTACT G

31

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13812

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GGCACCTCAA TGATCCTGGG TCAGC

25

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13804

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GCTGACCCAG GATCATTGAG GTGCC

25

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13815

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TGAGCTGGAT CCGCAGGAAG GGCAAACCA

29

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13811

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AGCTCAGGAT CCCTCCCAGG AGTGCAGG

28

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13810

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TGAGCTTCTA GATTAGCAGG AAGGGCAAAC

30

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13814

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AGAGCTGGAT CCCTCAGGAG CAGAGAAGAT CCC

33

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC13813

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AGAGCTTCTA GATTACTCAC GAGCAGAGAA GATC

34

CLAIMS

1. An isolated polynucleotide molecule selected from the group consisting of:

(a) polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO: 1 from nucleotide 97 to nucleotide 1215;

(b) orthologs of (a);

(c) allelic variants of (a) or (b);

(d) polynucleotide molecules that encode a polypeptide that is at least 80% identical to the amino acid sequence of SEQ ID NO: 2 from amino acid residue 22 (Leu) to amino acid residue 394 (Glu); and

(e) degenerate nucleotide sequences of (a), (b); (c) or (d).

2. The isolated polynucleotide molecule of claim 1, wherein the polynucleotide molecules comprise a nucleotide sequence as shown in SEQ ID NO: 1 from nucleotide 97 to nucleotide 1344, or polypeptide molecules that encode a polypeptide that is at least 80% identical to the amino acid sequence of SEQ ID NO: 2 from residues 22 to 437.

3. The isolated polynucleotide molecule of claim 1, wherein the polynucleotide molecules comprise a nucleotide sequence as shown in SEQ ID NO: 1 from nucleotide 79 to nucleotide 1344, or polypeptide molecules that encode a polypeptide that is at least 80% identical to the amino acid sequence of SEQ ID NO: 2 from residues 16-437.

4. The isolated polynucleotide molecule of claim 1, wherein the polynucleotide molecules comprise a nucleotide sequence as shown in SEQ ID NO: 1 from nucleotide 34 to nucleotide 1344, or polypeptide molecules that encode a polypeptide that is at least 80% identical to the amino acid sequence of SEQ ID NO: 2 from residues 1-347.

5. An isolated polynucleotide molecule comprising a sequence of nucleotides from nucleotide 97 to nucleotide 582 as shown in SEQ ID NO: 1.

6. An isolated polynucleotide molecule comprising a sequence of nucleotides from nucleotide 655 to nucleotide 1215 of SEQ ID NO: 1.

7. The isolated polynucleotide molecule of claim 6, wherein the polynucleotide molecules comprise a sequence of nucleotides from nucleotide 655 to nucleotide 1344 of SEQ ID NO: 1.

8. An isolated polynucleotide molecule encoding a fusion protein comprising a first polypeptide and a second polypeptide, said first polypeptide encoded by a sequence of nucleotides as shown in SEQ ID NO: 1 from nucleotide 97 to nucleotide 582, and said second polypeptide encoded by a sequence of nucleotides as shown in SEQ ID NO: 1 from nucleotide 655 to nucleotide 1215.

9. The isolated polynucleotide molecule of claim 1, wherein the polynucleotide is DNA.

10. An expression vector comprising the following operably linked elements:

a transcription promoter;

a DNA segment selected from the group consisting of:

(a) polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO: 1 from nucleotide 97 to nucleotide 1215;

(b) orthologs of (a);

(c) allelic variants of (a) or (b);

(d) polynucleotide molecules that encode a polypeptide that is at least 80% identical to the amino acid

sequence of SEQ ID NO: 2 from amino acid residue 22 (Leu) to amino acid residue 394 (Glu); and

(e) degenerate nucleotide sequences of (a), (b); (c) or (d); and a transcription terminator.

11. A cultured cell into which has been introduced an expression vector according to claim 10, wherein said cell expresses the polypeptide encoded by the DNA segment.

12. A method of producing a polypeptide comprising: culturing a cell into which has been introduced an expression vector according to claim 10, whereby said cell expresses a polypeptide encoded by the DNA segment; and recovering the polypeptide.

13. An isolated polypeptide selected from the group consisting of:

(a) polypeptide molecules comprising an amino acid sequence as shown in SEQ ID NO: 2 from amino acid residue 22 (Leu) to amino acid residue 394 (Glu);

(b) orthologs of (a);

(c) allelic variants of (a) or (b); and

(d) polypeptide molecules that are at least 80% identical to the amino acids of SEQ ID NO: 2 from amino acid residue 22 (Leu) to amino acid residue 394 (Glu).

14. The isolated polypeptide of claim 13, wherein said polypeptide molecules comprise an amino acid sequence as shown in SEQ ID NO: 2 from amino acid residue 22 (Leu) to amino acid residue 437 (Cys).

15. The isolated polypeptide of claim 13, wherein said polypeptide molecules comprise an amino acid sequence as shown in SEQ ID NO: 2 from amino acid residue 16 (Leu) to amino acid residue 437 (Cys).

16. The isolated polypeptide of claim 13, wherein said polypeptide molecules comprise an amino acid sequence as shown in SEQ ID NO: 2 from amino acid residue 1 (Met) to amino acid residue 437 (Cys).

17. An isolated polypeptide comprising a sequence of amino acids as shown in SEQ ID NO: 2 from amino acid residue 22 (Leu) to amino acid residue 183 (Pro).

18. An isolated polypeptide comprising a sequence of amino acids as shown in SEQ ID NO: 2 from amino acid residue 208 (Phe) to amino acid residue 394 (Glu).

19. The isolated polypeptide of claim 18, wherein said polypeptide comprises a sequence of amino acids as shown in SEQ ID NO: 2 from amino acid residue 208 (Phe) to amino acid residue 437 (Cys).

20. An isolated fusion protein comprising a first polypeptide and a second polypeptide, said first polypeptide comprising a sequence of amino acid residues as shown in SEQ ID NO: 2 from amino acid 22 (Leu) to amino acid 183 (Pro), and said second polypeptide comprising a sequence of amino acid residues as shown in SEQ ID NO: 2 from amino acid 208 (Phe) to amino acid 394 (Glu).

21. An isolated polypeptide according to claim 13, covalently linked at the N-terminus or the C-terminus to a moiety selected from the group consisting of affinity tags, toxins, radionucleotides, enzymes or fluorophores.

22. An isolated polypeptide according to claim 21, further comprising a proteolytic cleavage site between said sequence of amino acid residues and said affinity tag.

23. An antibody that specifically binds to an epitope of a polypeptide comprising a sequence of amino acids as shown in SEQ ID NO: 2 from amino acid 1 (Met) to amino acid 437 (Cys).

24. An oligonucleotide probe or primer comprising at least 14 contiguous nucleotides of polynucleotide of SEQ ID NO: 1 from nucleotide 34 to nucleotide 1344.

25. A pharmaceutical composition comprising a polypeptide according to claim 13, in combination with a pharmaceutically acceptable vehicle.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/09584

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/47 A61K38/16 C12N15/62 C07K16/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 32286 A (MICROGENESYS INC) 30 November 1995 see the claims see abstract; examples 1,2,4 see page 16, line 29 - page 17, line 14 ---	
A	WO 95 04548 A (JENNER TECHNOLOGIES) 16 February 1995 see the claims see abstract see page 8 - page 15 ---	
A	EP 0 747 705 A (BAYER AG) 11 December 1996 see the example see abstract ---	

-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 August 1998

Date of mailing of the international search report

04/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Oderwald, H

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/09584

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>NCI-CGAP: "Homo sapiens clone IMAGE:998299 (accession number AA535187)" EMBL SEQUENCE DATABASE, 26 July 1997, XP002075532 Heidelberg, Germany see the whole document -----</p>	24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/09584

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9532286	A	30-11-1995	AU 2517995 A	18-12-1995
WO 9504548	A	16-02-1995	AU 686660 B	12-02-1998
			AU 7631294 A	28-02-1995
			EP 0721345 A	17-07-1996
			JP 9504000 T	22-04-1997
EP 0747705	A	11-12-1996	JP 9005329 A	10-01-1997